



Utilisation of Green Manures as Organic Substrates in Soilless Culture: A Review

Elakiya, N. ^{a++*} and K.P. Ragupathi ^{b#}

^a J. K. K. Munirajah College of Agricultural Science (Affiliated to TNAU), Gobichettipalayam, Erode, Tamil Nadu, India.

^b College of Agricultural Technology, Gullapuram, Theni, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i72156>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118987>

Review Article

Received: 15/04/2024

Accepted: 20/06/2024

Published: 22/06/2024

ABSTRACT

Green manuring with inclusion of legumes is the most feasible option to fix atmospheric nitrogen up to 80-100 kg of nitrogen in 45-60 days. They are easily decomposable improving soil organic carbon, nutrient availability, physicochemical and biological properties of soil, and crop productivity and hence can be used as a substrate in the form of dry powder in soilless culture. The scope of utilizing organic materials for crop production remains viable in recent years particularly in containerized crop production. Soilless culture as a crop production system could solve global issues such as the shortage of water, environmental pollution and instability of ecological system in various ways providing optimum environmental medium for crop growth in order to gain maximum yield and high quality products. Henceforth, less land area is required for agriculture production

⁺⁺ Assistant Professor (Soil Science and Agricultural Chemistry);

[#] Assistant Professor (Seed Science & Technology);

*Corresponding author: E-mail: elakiya.nandu@gmail.com;

system resulting in increased land productivity. With this background, this review projects to maximize the utilization of the properties of green manures as an alternative grow-media substrate in the emerging soilless cultivation system.

Keywords: *Daincha; grow-media; legumes; sustainability.*

1. INTRODUCTION

“The world population is growing, and it is anticipated to reach 9.7 billion by 2050” [1,2]. Meeting demand will require a 50% increase in food supply by 2030 and a 70% increase in the next 40 years [3]. However, population expansion, industry, and urbanization are encroaching on arable land for food production, and it is expected to shrink by one-third of its current availability (Fig. 1) [4,5]. New food production technology is being developed to support agriculture's transition in response to population expansion and resource demands [6,7]. “Soilless culture is an emerging and promising technique for overcoming current agricultural threats. It is the most sustainable and environmentally acceptable alternative to traditional soil-based intensive agriculture, is now regarded as a convenient approach for conserving energy, fertilizers, and water. The biggest tasks handled in closed soilless systems are salinity of circulating solution, non-uniform nutrient supply, root pathogen control, removal of undesired taste and ill-health related compounds in the product, and so forth” [8].

“Green manuring has become a very important agricultural practice since last three decades, and green manuring practices are promising tools for the better sustainability of agriculture in the longer run and with almost no hazards to the environment” [9]. “Green manures (GM) rich in N and P readily decompose, hence can also be used as a substrate in the form of dry powder for

grow-media culture” [10]. Hence, this review gives knowledge regarding the composition, rates and placement of GM crops, in future, to propose holistic system and precise agricultural practices to deal with constrain during adaptation of GM-based cropping system and utilization of green manure dry powders as one of the choice as alternative substrate for soilless cultures.

2. UTILIZATION OF ORGANIC SUBSTRATES FOR SOILLESS CULTURE

Soilless culture systems are currently taking up new dimensions in the modern greenhouse industry due to their several advantages. Use of recyclable organic resources as soilless media is gaining importance worldwide during last fifty years [11] for intensive plant production due to environmentally friendly disposition [12] of resources. “The term ‘soilless culture’ generally refers to any method of growing plants without the use of soil as a rooting medium” [13]. “Soilless culture is a distinct technology used to produce agricultural and horticultural crops” [14]. It entails growing crops in an absence of soil in a closed environment, protected cultivation, and vertical farming. The most accurate control over the supply of water, nutrients, pH, temperature, etc. are the major advantages of soilless culture which apparently increase the crop productivity by reduction of labour and making more crops per year. The common soilless culture only takes eight to ten weeks for crop harvesting.




Fig.1. World population (billions) versus arable land (ha per person) 1950-2050 [15]

The soilless crop production system is divided into two categories: (i) water-based culture, such as deep water culture (DWC by Gericke 1929), float hydroponics (developed by University of Arizona, 1860), nutrient film technique (NFT, Allen Copper, 1965), and aeroponics (suggested by Richard Stoner, 1983) [16] and (ii) substrate-based culture, or substrate culture using nutrient/substrate growing media. Soilless systems enable optimization of both physical and chemical characteristics in the root environment and a more efficient control of pathogens. As a result, higher yields at a reasonable production cost can be attained [17].

“Growing media have been used for both the production of high-value vegetables and ornamental plants, as well as for plant propagation. Growing media or substrates are defined as all solid materials, other than soil, which either alone or in mixtures can guarantee better plant growth conditions than agricultural soil in one or many aspects” [18]. “Organic constituents are majorly used in soilless media composition. Need based utilization of different inorganic substrates is made in growing media which include rockwool, perlite, pumice, zeolite, tuff, volcanic porous rock, expanded clay granules, and vermiculite” [13].

Organic substrates can be synthetic or can consist of a natural organic matter. Substrate culture uses natural organic nutrient substrates such as peat, coir, plant waste, sawdust, bark, rice hulls, rice husk, rice straw, compost, vermicompost, meal, cake, farm yard manure (FYM), coco peat, brick shards, bio control agents, bio fertilizers, paper waste, wood sawdust, peat moss, sphagnum moss, bagasse or natural substrates such as sand, gravel, tuff, pumice, perlite, rock wool, vermiculite, montmorillonite, etc. as the growing medium.

“The most available and applicable organic materials are peat, composts, bark and wood residues. The swamp rose mallow grown in containers with vermicompost showed improved plant dry weight” [19]. “Vermicompost mixed with coir in 2:1 ratio as container substrate increased plant height and fresh weight in *Beta vulgaris* L”. [20]. Bhat et al. [21] revealed that “substrates containing vermicompost, coco-peat, perlite and sphagnum peat moss (2:1:1:1 or 1:1:1:1 v/v) produced significantly better growth, yield and quality in tomato, cucumber and capsicum than other substrate combinations”. Kameswari et al. [22] obtained “highest values of plant height,

plant spread, number of branches per plant, duration of flowering, flower weight, spray length and number of flowers per plant were in the potting media containing cocopeat + sand + FYM+ vermicompost”.

“Different substrates were compared for soilless culture in watermelon and the plant growth, yield, fruit quality and plant nutrient uptake were highest with vegetative growth in basaltic mix, sand, peat and greenhouse soil respectively and highest yield was in perlite” [23]. Majdi et al. [24] recorded “peat + perlite had most effect on growing traits and yield of green pepper”. Huang et al. [25] studied “the combined effect of biochar and vermicompost amendments on the plant growth of basil and tomato. The results showed that plants in Biochar: Vermicompost: Peat based substrate exhibited similar growth indexes and total dry weight as in 100 % Peat based substrate”.

2.1 Scope of Utilizing Organic Substrates in Soilless Crop Production

The stability of soil organic matter depends on the sustained addition of organic materials in soils by way of incorporation of crop residues and organic manures and on the rightly driven biological activity. Formation of particulate organic matter in accountable proportion imparts major roles in continuous cropping systems where integrated nutrient management practices are judiciously undertaken. Prevailing conducive agro-climatic conditions also favour more accumulation of particulate organic matter. Recycling of organic matter in soils is a long-term process.

The scope of utilizing organic materials for crop production remains viable in recent years particularly in containerized crop production. Finding proper combination of organic materials to compose effective growing media for anchoring root and supporting plants adequately with water and nutrients is the need of the hour.

3. GREEN MANURING CROPS

Green manuring is the action of agronomic practices and assimilation of legume and non-legume green plants into the soil either by adopting in-situ or the plants developed abroad and congenial into the soil for abundance improvement. Green manures are made by incorporating green plant

materials (either freshly cut weeds or rotation crop debris) into soils. Green admixture crops may include legume crops; non-grain, aroma, and beat legumes; abiding coarse multifunctional shrubs; and crops suitable for green leaf manuring (Table 1). "Upon decay, green manures enrich the soil with organic matter and to a lesser extent with nutrients such as nitrogen and phosphorus. Ample time for the green manure to decompose is allowed between the incorporation of

weed/crop debris in the soil and the planting of the new crop" [26].

Legume crops and tree species suitable for green manuring in different agro-climatic zones are mentioned in Table 2 and nutrient composition of some of the green manures and green leaf manures are given in Fig. 2. and some of common green manures and green leaf manures of tropical region are depicted in Fig. 3. and Fig. 4.




Fig. 2. Nutrient content of green manures

Table 1. Categories of green admixture crops [27]

Legume crops	Non-grain, aroma and beat legumes	
Pigeon pea (<i>Cajanus cajan</i>)	Sunn hemp (<i>Crotalaria juncea</i>)	Subclover (<i>Trifolium subterraneum</i>)
Green gram (<i>Vigna radiata</i>)	Dhaincha (<i>Sesbania species, S. Aculeata</i>)	Strawberry clover (<i>T. Fragiferum</i>)
Cowpea (<i>Vigna unguiculata</i>)	Wild indigo (<i>Tephrosia purpurea</i>)	Persian clover (<i>Trifolium resupinatum</i>)
Soybean (<i>Glycine max</i>)	Black henna (<i>Indigofera tinctoria</i>)	Red clover (<i>Trifolium pratense</i>)
Groundnut (<i>Arachis hypogea</i>)	Barseem (<i>Trifolium alexandrinum</i>)	White clover (<i>Trifolium repens</i>)
Cluster bean (<i>Cyamopsis tetragonoloba</i>)	Broadbean (<i>Vicia faba</i>)	<i>Desmodium</i>
Sub-tropical grasses and weeds	White lupin (<i>Lupinus albus</i>)	<i>Centosemia</i>
<i>Panicum maximum</i>	Blue lupin (<i>Lupinus angustifolius</i>)	<i>Stylosanthes</i>
<i>Pennisetum purpureum</i>	Yellow lupin (<i>Lupinus luteus</i>)	
<i>Tripsacumlaxum</i>	Common vetch (<i>Vicia sativa</i>)	
<i>Aduthoda vesica</i>	Fenugreek (<i>Trigonella foenumgraecum</i>)	
<i>Eicchornia crassipes</i>	Candied clover (<i>Melilotus spp.</i>)	
<i>Trianthema portulacastrum</i>	Trefoil (<i>Lotus spp.</i>)	
<i>Ipomoea carnea</i>	Black medic (<i>Medicago lupulina</i>)	
<i>Calotropis gigantea</i>	Lucerne or alfalfa (<i>Medicago satvia</i>)	
Green Leaf manures		
<i>Leucaena leucocephala</i> (Subabul)	<i>Azadirachta indica</i> (neem)	<i>Tephorsia candida</i>
<i>Gliricidia spp.</i>	<i>Derris indica</i>	<i>Dodonea viscosa</i>
<i>Cassia siamea</i> (Kassod tree)	<i>Cassia tora</i>	<i>Hibiscus viscosa</i>
<i>Cassia auriculata</i>	<i>Cassia accidentalis</i>	<i>Delonix elata</i>
<i>Delonix regia</i>	<i>Peltophorum ferrugenum</i>	<i>Cassia nigricans</i>

Table 2. Leguminous green manure crops for different regions [28,29,30]

(A) In situ green manure crops			
(a) Tropical region		(b) Temperate region	
Common name	Scientific name	Common name	Scientific name
Cluster bean	<i>Cyamopsis tetragonoloba</i>	Subterranean clover	<i>Trifolium subterraneum</i>
Cowpea	<i>Vigna unguiculata</i>	Ladino clover	<i>Trifolium repens</i>
Pueraria	<i>Pueraria phaseoloides</i>	Crimson clover	<i>Trifolium incarnatum</i>
Green gram	<i>Vigna radiata</i>	Faba bean	<i>Vicia faba</i>
Lablab	<i>Lablab purpureus</i>	Soybean	<i>Glycine max</i>
Dhaincha	<i>Sesbania aculeate, S. rostrata</i>	Red clover	<i>Trifolium pratense</i>
White lupin	<i>Lupinus albus</i>	Black lentil	<i>Lens culinaris</i>
Gray bean	<i>Mucuna cinerecum</i>	Alfalfa	<i>Medicago sativa</i>
Pigeon pea	<i>Cajanus cajan</i>	Barrel medic	<i>Medicago truncatula</i>
Sunn hemp	<i>Crotalaria breviflora, C. juncea, C. striata</i>	Hairy vetch	<i>Vicia villosa</i>
Buffalo bean	<i>Mucuna aterrima</i>	Milk vetch	<i>Astragalus sinicus</i>
Jack bean	<i>Canavalia ensiformis</i>	Winter pea	<i>Pisum sativum</i>
Velvet bean	<i>Mucuna deeringiana</i>	Sweet clover	<i>Melilotus officinalis</i>
Stylo	<i>Stylosanthes guianensis</i>	Cura clover	<i>Trifolium ambiguum</i>
Desmodium	<i>Desmodium ovalifolium</i>	Purple vetch	<i>Vicia benghalensis</i>
Milk vetch	<i>Astragalus sinicus</i>	Common vetch	<i>Vicia sativa</i>
Zornia	<i>Zornia latifolia</i>	(B) Ex situ green leaf manuring shrubs and trees	
Jumby bean	<i>Leucaena leucocephala</i>	Common name	Scientific name
Kudzu	<i>Pueraria phaseoloides</i>	Subabul	<i>Leucaena leucocephala</i>
Adzuki bean	<i>Vigna angularis</i>	<i>Vigna angularis</i>	<i>Gliricidia</i>
Black gram	<i>Phaseolus mungo, P. trilobus</i>	<i>Phaseolus mungo, P. trilobus</i>	<i>Gliricidia sepium</i>
Soybean	<i>Glycine max</i>	<i>Glycine max</i>	<i>Pongamia glabra</i>
Alfalfa	<i>Medicago sativa</i>	<i>Medicago sativa</i>	<i>Calotropis gigantea</i>
Wild indigo	<i>Indigofera tinctoria</i>	<i>Indigofera tinctoria</i>	<i>Tephrosia purpurea</i>
Berseem	<i>Trifolium alexandrinum</i>	<i>Trifolium alexandrinum</i>	<i>Indigofera teysmannii</i>
			<i>Sesbania speciosa, S. rostrata</i>
			<i>Cassia tora</i>



Cluster beans
Cyamopsis tetragonoloba



Cowpea
Vigna unguiculata



Pueraria
Pueraria phaseoloides



Green gram
Pueraria phaseoloides



Lablab
Lablab purpureus



Dhaincha
Sesbania aculeate, S. rostrata

Fig. 3. Green manures of tropical region



White lupin
Lupinus albus



Berseem
Trifolium alexandrinum



Pigeon pea
Cajanus cajan



Sunn hemp
Crotalaria breviflora, *C. juncea*, *C. striata*



Buffalo bean
Mucuna aterrima



Jack bean
Canavalia ensiformis

Fig. 3. Green manures of tropical region (contd.)



Velvet bean
Mucuna deeringiana



Stylo
Stylosanthes guianensis



Desmodium
Desmodium ovalifolium



Milk vetch
Astragalus sinicus



Zornia
Zornia latifolia



Jumby bean
Leucaena leucocephala

Fig. 3. Green manures of tropical region (contd.)



Kudzu
Pueraria phaseoloides



Adzuki bean
Vigna angularis



Black gram
Phaseolus mungo, P. trilobus



Alfalfa
Medicago sativa



Soybean
Glycine max



Wild indigo
Indigofera tinctoria

Fig. 3. Green manures of tropical region (contd.)



Gliricidia
Gliricidia sepium



Karanj
Pongamia glabra



Milkweed
Calotropis gigantea



Tephrosia
Tephrosia purpurea



Wild indigo
Indigofera teysmannii



Kassod
Cassia tora

Fig. 4. Green leaf manures of tropical region

3.1 Nitrogen Contribution by Green Manuring

“Nitrogen is the primary macronutrient responsible for plant vegetative development. Plants contain 1-6% nitrogen by weight, which is necessary for a number of plant metabolic processes. Plants consume it in the forms of NO_3^- (nitrate) and NH_4^+ (ammonium), as well as amide. It is a component of many metabolically active chemicals, including amino acids, protein, nucleic acids, purines, compounds, co-catalysts,

and alkaloids. Nitrogen in plants undergoes a few compound-catalyzed processes, resulting in NH_3 , which is absorbed into amino acids and subsequently consolidated into proteins and nucleic acids. It is an essential component of the porphyrin ring in chlorophyll, which turns light energy into chemical energy during photosynthesis. The rate of nitrogen uptake by plants at different developmental stages has a significant impact on yield and yield-attributing characteristics” [31].

“Green manures, particularly legumes, contribute a higher amount of nitrogen. They fix atmospheric di-nitrogen by symbiosis with rhizobia (*Rhizobium*, *Bradyrhizobium*, *Sinorhizobium*, and *Mesorhizobium*). The legume rhizobium symbiosis compensates for 40% of global fixed nitrogen” [32]. Fig. 5. depicts soil enrichment with nitrogen by fixation and mineralization. “Nitrogen supply in green manures varies depending on the species and total biomass generated” [33]. “Leguminous green manures have been reported to fix up to 80-100 kg of nitrogen in 45-60 days” [34]. “The *Azolla-Anabaena* symbiosis may fix 2-4 kg N/ha per day” [35]. “Green manure nitrogen is superior to urea nitrogen in terms of plant growth and development, as well as crop grain output” [36]

Nitrogen supply by green manures is not exigent in nature, it’s a gradual and slow process. Green manuring crops not only fix atmosphere di-nitrogen, but they also help to conserve nitrogen in the soil. By green manuring, the decomposing plant provides necessary nutrients for crop growth, reducing the need for chemical fertiliser

throughout the juvenile period. Green manures improve soil qualitative characteristics with comparable efficiency to mineral fertilisers, but mineral fertilisers deteriorate soil conditions [38]. However, concrete information on rate of decomposition and quantity of available of nutrients released is scanty with daincha incorporation [39].

A slower rate of nitrogen release by green manures reduces nitrogen losses by leaching, denitrification, and NH_3 volatilization. Soil supplemented with green manure showed no NH_3 volatilization losses. The portion of green-manure nitrogen available for a subsequent harvest is typically between 40% and 60% of the total amount contained in green manuring crops. Green manure nitrogen is divided into two parts: 'fast N' and 'slow N'. Fast N degrades quickly and is readily available to plants, whereas Slow N degrades slowly and is available to plants for an extended period of time [40]. Some green manuring crops along with their nitrogen contribution is presented in (Table 3).




Fig. 5. Soil enrichment with N by fixation and mineralization [37]

Table 3. N accumulation in major leguminous green manure crops

Crop species	Growth duration (days)	N accumulation (kg ha ⁻¹)	Reference
<i>Glycine max</i>	45	115	[41]
<i>Crotalaria juncea</i>	45	169	
<i>Cajanus cajan</i>	45	33	
<i>Sesbania aculeata</i>	45	225	
<i>Vigna radiata</i>	45	75	
<i>Dolichos lablab</i>	45	63	
<i>Indigofera tinctoria</i>	45	45	
<i>Sesbania rostrata</i>	56	176	[42]
<i>Sesbania aculeata</i>	56	144	
<i>Vigna unguiculata</i>	45	75	
<i>Vigna radiata</i>	45	75	[43]
<i>Sesbania rostrata</i>	60	219	[44]
<i>Sesbania cannabina</i>	60	171	
<i>Sesbania aegyptiaca</i>	57	39	[45]
<i>Sesbania grandiflora</i>	57	24	
Cluster bean	49	91	[36]
Common vetch	Flowering	105–210	
Sweet clover	Flowering	150–300	
Milk vetch	Flowering	65–131	[46]

Green manure crops can provide the rice crop with 50–80 kg N/ha [47]. According to Mann et al. [48], after three years of continuous green manuring, soil organic matter, N, and P grew to 1.09%, 0.37%, and 10.2 ppm respectively. Saravana and Perumal [49] concluded that the addition of green manure (*S. aculeata*) and fertiliser N together enhanced the status of accessible N. *S. rostrata* had the highest total soil nitrogen levels, followed by *S. aculeata*, and green gram incorporation had the lowest levels [50].

The soil nutrient balance after incorporation of different green manures specially *S. rostrata*, *S.*

aculeata and *C. juncea* exhibited positive balance of nutrients than other green manures [52]. While evaluating the effect of different in-situ green manuring on soil organic matter (%), N (%), P (ppm) and K (meq/100 gm) contribution on pre-sown rice soil in 2015 and 2016, Irin and Biswas [51] concluded that after two consecutive year, Green Manure-Transplant Aman-Mustard Cropping Pattern increased soil organic matter, nitrogen, phosphorus and potassium compared to initial soil by which N fertilizer rates could be reduced after the incorporation of green manures in the succeeding and following crops (Fig. 6).




Fig. 6. Effect of in situ green manure incorporation on soil properties at two consecutive years (T0= Control, T1= *S. aculeata*, T2= *S. rostrata*, T3= *C. juncea*, T4= *V. radiata*, T5= *V. mungo*, T6= *V. unguiculata*, T7= *L. leucocephala*, T8= *M. pudica*) [51]

3.2 Contribution of Other Nutrients by Green Manuring

Green manures, especially those with deep roots, collect nutrients from deeper regions and make them available after decomposition. Green manuring crops release and recycle nutrients (nitrogen, phosphorus, and potassium) during decomposition, which aids in integrated plant nutrient management [53,54,55,56,57,58,59].

“Phosphorus availability is frequently reduced in calcareous and acidic soils due to interaction with calcium carbonate and iron oxide” [60]. “Phosphorus release following decomposition is typically associated with phosphorus levels in green manure. About 40 to 60 percent of phosphorus is immediately released during the degradation of plant biomass. In organically managed system, mineralization of available organic phosphorus in soil is the primary source of phosphorus. The phosphorus in the soil has to be made available to the plants” [61]. “Green manure absorption into soil improves the phosphorus cycle and increases the availability of sparingly soluble phosphorus” [62,63]. “Green manure crops collect a considerable amount of P, which decomposes into bicarbonates (H_2CO_3). This bicarbonate can solubilize soil mineral P, ensuring sufficient phosphorus for subsequent crops” [64,65]

“Addition of green manures increase the soil organic carbon subsequently leading forward to reduction on soil pH. This decrease in soil pH reduces the phosphate fixation in soil with iron and aluminium” [66]. “Ultimately availability of phosphorus increases. Lupins grown in phosphorus deficient soil were found to extrude protons and different organic acids” [67,68]. “Green manuring uplifts the P uptake of succeeding crops by converting the fixed phosphorus into readily available forms” [62]. “P deficiency stimulates the formation of cluster roots in green manuring crops which are more active in P mobility and uptake” [68]. “In waterlogged conditions, green manures increased the availability of P through the mechanism of reduction, chelation and favourable alteration in soil pH. Higher availability of phosphorus from rock phosphate was reported due to green manuring in the rice fields” [62].

“On decomposition, organic P in green manure biomass could provide a relatively labile form of P to succeeding crops, thereby create a larger

pool of mineralizable soil organic P to supplement soluble inorganic P pools. The extensive root system of green manure crops improved the physical condition of the soil and liberated CO_2 and organic acids, which help in dissolving native potassium in soil and thereby increases the availability of potassium and per cent increase noticed was 2.1 to 4.9 %” [70]. Incorporation of *Sesbania aculeata* and *Crotalaria juncea* with N significantly increased soil organic carbon, N, P, K, Ca, and S [69] (Table 4).

“Green manure crops can fix up to 153 kg K/ha and up to 20 kg P/ha” [71]. “Dhaincha and green gram increased the soil available potassium by 3.7 and 2.4 per cent respectively. P and K utilization to an extent of 10 to 12 per cent was observed in field conditions due to green manure incorporation. GM crops contain appreciable amounts of NPK including other trace elements also” [72]. “They also mobilize S, P, Si, Zn, Cu, Mn and other nutrient element as a result of increased microbial activity (CO_2 formation) and decreased redox potential” [73]. “Green manuring with *Sesbania rostrata* increased both availability in soil and accumulation in plant of Fe, Mn and Cu due to the development of intense reducing condition, complex formation and greater nutrient holding capacity” [74]. Eriksen and Thorup-Kristensen [75] found that cruciferous crops such as winter rape or fodder radish were particularly effective at preventing sulphur leaching into lower soil profiles. Green manures such as forage chicory accumulate large amounts of micronutrients including sulphur, boron, manganese, molybdenum, and zinc [76]. Green manuring crops promote mycorrhizal growth on the roots of succeeding crops, increasing soil phosphorus (P) and micronutrient availability [77]. Soil P and K contents were highest in plot with *Crotalaria*, while soil Ca content was highest in plot with *Mucuna* [78].

Decomposition of daincha biomass increases humus, available nitrogen and lower down the C:N ratio of soil. Nutrient content of 60 days old dhaincha plants had 3 per cent N in addition to K, Ca, Mg, P, S and micronutrients. Addition of dhaincha biomass to soil at $1.0 t ha^{-1}$ added about 33 kg N, 1 kg P, 14 kg K, 14 kg Ca, 16 kg Mg and 2 kg S [79]. Across maize developmental stages, the green manuring treatment increased dry matter accumulation and N uptake by 28-114% and 83% to 146%, respectively. Green manure treatment significantly increased soil

Table 4. Effect of green manuring crops and nitrogen levels on physical and chemical properties of postharvest rice soil [69]

Treatment	Bulk density (g/cc)	pH	OC (%)	Total N (%)	Available P (mg/kg)	Exchangeable K (meq/100 g soil)	Available S (mg/kg)
Initial value	1.35	5.76	0.81	0.064	4.86	0.138	21.21
Control	1.32	5.73	0.83	0.055	4.69	0.129	20.57
<i>S. aculeata</i> + N0	1.19	5.60	0.90	0.073	5.75	0.205	22.38
<i>S. aculeata</i> + N15	1.17	5.60	0.92	0.081	6.11	0.219	22.61
<i>S. aculeata</i> + N30	1.15	5.60	0.93	0.093	6.23	0.223	22.83
<i>S. aculeata</i> + N45	1.13	5.61	0.93	0.098	6.43	0.248	23.01
<i>S. aculeata</i> + N60	1.11	5.62	0.97	0.106	6.71	0.269	23.19
<i>C. juncea</i> + N0	1.19	5.61	0.89	0.069	5.86	0.194	22.53
<i>C. juncea</i> + N15	1.19	5.61	0.89	0.075	6.16	0.216	22.76
<i>C. juncea</i> + N30	1.17	5.62	0.90	0.088	6.38	0.219	22.91
<i>C. juncea</i> + N45	1.15	5.63	0.91	0.095	6.55	0.242	23.13
<i>C. juncea</i> + N60	1.12	5.62	0.94	0.103	6.83	0.261	23.36
N60	1.22	5.67	0.84	0.083	5.58	0.248	22.15
CV (%)	2.25	2.01	2.55	2.05	4.95	3.060	2.23
LSD (0.05)	0.04	0.19	0.039	0.003	0.51	0.011	0.28

organic carbon by 3.90–12.23% over all N application rates, and total nitrogen and available nitrogen were significantly increased by 3.79–15.76% and 4.87–17.29%, with total phosphorus and available phosphorus by 6.1–13.6% and 9.6–5.3% respectively during maize developmental stages in the North China Plain [80].

In green manuring plot, the number of filled grain/panicle and highest grain yield (4t/ha) was obtained due to expected slow released nitrogen available throughout the growth period of rice [81]. After three years of continuous green manuring, the soil organic matter and N increased up to 1.09% and 0.37%, respectively [82]. Accountable increase in soil organic matter depended on the biomass producing ability of different dhaincha accessions [83]. Root, leaf and stubbles of dhaincha after decomposition improved the organic matter status of soil on green manuring due to that organic matter and total N status of soil ranged from 1.42 to 1.58% from initial level 1.51% and from 0.075 to 0.098% from initial level 0.078 per cent respectively after three years of continuous incorporation [84].

When comparing the different natural farming practices on yield of black gram, combination of Beejamrutha, Jeevamrutha, mulching and green manure treatment significantly recorded higher growth and yield attributes along with yield (1.062 t/ha) [85] which may be attributable to higher nutrient availability throughout crop growth, which was further ensured by improved microbial activity in the soil [86,87,88]. In rice-based cropping system, the rice grain yield increased 32 to 77 per cent over control due to incorporation of daincha as green manure along with different doses of NPK fertilizers, which can be attributed to the efficient and adequate nutrients supply from daincha biomass decomposition and release nutrients for utility of the crop [89,90].

Among the organic manures, *Sesbania aculeata* played a vital role in improving the uptake of NPK due to quick release of N from the added green manure with increased availability of P through the mechanism of reduction and chelation at favourably changed soil pH, release of K through the priming effect and direct contribution of K by green manure [91]. Effect of incorporation of dhaincha biomass in soil was noticed in increased organic matter content and total nitrogen. The amount of organic matter content varied from 1.582 to 2.133 per cent before

incorporation and 1.995 to 2.271 per cent after incorporation of dhaincha biomass in soil. Likewise, the total N in soil varied from 0.088 to 0.118 per cent before incorporation and 0.106 to 0.126 after incorporation [92].

Application of 100 percent NPK through inorganic fertilizers along with 6.25 t ha⁻¹ dhaincha found out from resulted in higher nutrient uptake in crop, greater soil available nitrogen, phosphorus, potassium and organic carbon in soil. This effect was followed in application of 100% N through dhaincha + balance P and K through inorganic fertilizers [93]. Green manuring enhanced boron and iron content of soil. Further, decomposed materials of *Sesbania* also serve as chelating compounds and help in increasing the availability of metallic micronutrients like Zn, Cu, Mn in succeeding crop [37]. Incorporation of dhaincha remarkably increased soil pH, organic carbon, N, P, K and S compared to initial soil nutrients status at the peak period of 50 days after incorporation and declined thereafter except in soil pH, K and S [94].

3.3 Improvement in Soil Conditions on Green Manuring

Daincha incorporation as green manure improved the soil physical environment, made the soil softer indicated by reduced bulk density, increased porosity of soil, increased the availability of major nutrients and ultimately favoured in increased yield of rice [95]. Further, green manuring with daincha crop improves soil structure, aeration, permeability and also protect leaching of nutrients from the soil. Green manuring effects on crop growth and nutrient utilization are associated with an improvement in soil physiochemical properties, such as bulk density, water conductivity, and carbon and N levels [96,97].

Soil water stored in 2-year *Sesbania* improved fallows was greater than in continuously cropped fertilized or unfertilized maize in eastern Zambia [98]. Incorporation of green manures reduced soil bulk density compared with the control and NPK fertilizer in consecutive years (Fig.7). However, there were no significant differences in bulk density between Moringa, Pawpaw, Mesquite and Neem leaves used as green manure [99]. Water-stable aggregate ratio (%) and carbon percent of soil macro-aggregates (>250 µm) and micro-aggregates (53 - 250 µm) in green manure treated plot were significantly higher than in NPK

(Fig. 8). Thus, green manure promoted soil aggregation and stabilized carbon in soil aggregates confirming green manure supplies nutrient, improves soil carbon storage and soil physical stability [100].

Incorporation of green manures had slightly lowered soil pH compared with control and this decrease in soil pH may be due to production of CO₂ and organic acids during decomposition of incorporated green manures [69] (Table 4).

Reduction from initial soil pH (7.64 to 7.96) was observed after green manuring resulting to lower pH (7.47 to 7.88). The reduction in pH might be due to incorporated *Sesbania* leaves whose sap has a pH of 4.0 and strongly acidic in nature. After decomposition, a marked influence in neutralizing the high pH of soil occurred due to production of organic acids [70,99]. Some other authors also reported that application of different types of green manures also decreased soil pH [101,102].




Fig. 7. Effect of various green manures and NPK fertilizer on soil bulk density [99]




Fig. 8. Distribution of water-stable aggregate and carbon of each fraction in treatments: CN, control; NPK, chemical fertilizer treatment; GM, green manure treatment [100]

Salinity of the soil increases after adding green manure due to increase in cations and anions or increasing in release of salts and nutrients from decomposing biomass. Organic carbon content significantly improved by incorporation of *Sesbania* green manure upto 6.7 to 11.8 per cent over check. The status of available N was improved when *Sesbania* was incorporation at 45 days of growth from 2.7 per cent in check plots to 4.8 percent in daincha incorporated plots [70].

Soil available P extracted was highest in green manured fields which may be due to the improved physical condition of the soil, by providing aeration and enhancing the microbial activity and mineralization of P from green manure biomass as well as chelation of calcium in alkaline soil and aluminium in acid soils [103]. Incorporation of daincha at 45 days of growth resulted in slight increase in available phosphorous from 21.2 to 22.3 kg ha⁻¹ [70].

3.4 Effect of Green Manuring on Biological Activity

The decomposition of green manures serves major functions for microflora providing both C and energy for growth and formation of new cell material, which further multiplies its colony saprophytically on the decomposing OM [104]. A large number of soil microorganisms exist in the soil as long as there is a C source for energy [9]. Green manures promote soil microbial growth and activity, mineralization [105,106], and improve soil fertility and quality [107]. Soil microbes decompose organic matter by producing numerous enzymes, which increases soil enzymatic activity [53,108]. Green manure provides nutrients rich in organic carbon to the microbial biomass and increases the biodiversity of soil microorganisms, resulting in disease suppression, and improved soil structure, soil properties, and crop health [9,109,110].

“GM has two main positive points from the microbiological point of view: (a) primarily it provides nutrient-rich OM for the microbial community which easily converts organically bound nutrients in plant residues to easily available nutrient form to the crops; (b) secondly it enhances the biodiversity of soil microorganisms. This microbial diversity can be increased by incorporating different legume green manure in crop rotation and cropping system programs” [105,110].

Green manure treatments can significantly increase the number of soil fungus, bacteria, and actinomycetes [111]. “Green manure treatment

increased soil microbial biomass carbon and nitrogen by 1.94%-93.07% and 2.30%-145.07%, respectively” [104]. “Soil microbial biomass carbon and nitrogen is an important index for assessing rhizosphere effect” [112]. Green manures can supply nutrients for the reproduction of soil microbes, increasing diversity and making the environment suitable for new microorganisms to develop in that site. Green manure applications not only enhanced the amount of soil microorganisms, but also boosted soil enzymatic activities due to root exudations and the efficiency of soil nutrient transformation, which is all advantageous.

Due to soft and succulent nature, dhaincha plants decompose readily in soil within 60 days. The rate of decomposition is very rapid within one to two weeks after incorporation and then decline gradually [113]. Daincha fixes atmospheric nitrogen through legume-Rhizobium symbiosis and hence its accumulated nitrogen in above ground matter has the potentiality to enhance growth and yield of subsequent crops when incorporated and decomposed [114,115]. The increase in total N content of soil due to incorporation of daincha may be attributed to the mineralization of N green manure biomass due to greater multiplication of soil microbes, which could have converted organically bound N to inorganic form [8] in rice-based cropping system [89].

“Mineralization of nutrients from decomposing green manure was initially very low later increased at a faster rate up to 5-7 weeks and thereafter declined with time” [84]. “Green manure decay rates would increase when N contents were increased or would decrease if the N content declined” [116]. “In wetland rice fields, 40 per cent of carbon and 80 per cent nitrogen of *Sesbania* mineralized in initial 10 days of incorporation while release of N was highest in the fifth week after incorporation” [117]. In another study, the peak N release occurred after 5-8 weeks of decomposition of green manure and declined thereafter. Decomposition rate becoming slow or declining after a certain period might be due to declined microbial activity controlled by soil temperature, oxygen, substrate availability and microorganism activity as well C:N ratio [118].

Green manure application, particularly when combined with organic fertilisers, increased alpha diversity in the soil bacterial community, whereas the fungus community showed the opposite tendency. It also changed the soil microbial

populations during both the growth and incorporation phases, particularly the taxa involved in the carbon, nitrogen, and sulphur cycles. Compared to fallow treatments, green manure application significantly raised microbial biomass carbon by 29.8%-72.9%, regardless of the kind of combination fertiliser used. Green manure increased urease activity by 35.6%-142.6% and β -glucosidase activity by 65.9%-172.9% compared to fallow treatments. However, it decreased catalase activity by 22.5%-55.6% [119] (Fig. 9).

“The mineralization rate of incorporated green manure was higher during the first week of its incorporation and extended till 20 days, then the process slowed down” [120]. “In the initial process of decomposition of Sesbania plant, the rate of decomposition is high and decomposable substances are released rapidly” [39]. Firstly,

celluloses are decomposed and used by microorganisms, more recalcitrant such as lignin and tannin are decomposed consequently decomposition rates decreased [121].

“In the initial stages of incorporation of daincha, leaf, bark and other soft parts break down and these parts contain small soluble carbon molecules like starches and amino acids which are easy to break down. However, more recalcitrant molecules like lignin consist of complex molecules; consequently, it takes a longer time of decomposition and thus nutrients reserve in soil and release latter” [122]. At 50 days after incorporation, the available nutrients were gradually declined possibly due to leaching of nutrients or fixation in soil. Sesbania as a legume provides high quality residues with lower C/N, lignin, and polyphenol contents [123].




Fig. 9. Effects of application of green manure combined with different fertilizers on (A) the soil microbial carbon, (B) urease activity, (C) β -glucosidase activity and (D) catalase activity during the green manure incorporation period. Control (CK) - gray bar, fallow treatments - green, ryegrass incorporation treatments - blue and Chinese milk vetch incorporation treatments - orange bars [119]

Ash, crude protein and crude fibre content in dhaincha contribute to increase the soil nutrients status [94,124]. "Histidine, arginine and aspartate are the most abundant amino acids produced in seed exudates of *Sesbania*. A number of bacteria can use amino acids such as arginine, histidine, glutamate, glutamine, alanine and alanine and other amino acids as a sole nitrogen source in the presence of a suitable carbon source" [125]. Chanda et al., [126] reported "biochemical properties of *Sesbania* species indicating substantial content of ash (9.03 %), crude fibre (25.57 %), crude protein (18.22 %), lignin (21.97 %), holo-cellulose (70.28 %) and alpha-cellulose (41.22 %)". "During four years of continuous experiment with ryegrass (*Lolium multiflorum* L.) application to soil, soil microbial biomass carbon and nitrogen, soil respiration, soil enzymatic activities such as urease, invertase, and catalase were improved when compared to the control" [104].

3.5 Potentials of Green Manure as a Substrate and Amendment

Green manures are also grown for animal fodder [127], ground cover, firewood and other uses in traditional agro-forestry systems. To minimise chemical inputs in agriculture, green manure has been adopted preference over inorganic fertiliser [96,128]. Daincha is a popular and ideal green manure because of quick growing nature and its green matter is succulent and easily decomposable at low moisture content. On addition to soil, daincha can add significant amount of organic matter and nitrogen [129].

Advantageously many green manure crops grow well even in marginal lands with little or no input. Daincha can exhibit luxuriant growth in soil with a high salinity up to electrical conductivity of 10 dS m⁻¹. Due to this stability daincha and other few *Sesbania* spp. have been recommended for reclamation of saline and sodic soils [130]. Long duration cultivation of dhaincha for about 10-12 year would eliminate desertification of marginal lands and hence its cultivation is recommended to rehabilitate degraded lands into productive crop lands [131]. Daincha crop has an annual yield potential of upto 20 tonnes of drymatter per hectare under appropriate production strategy [132]. At an dry matter production rate of 5.2 t ha⁻¹ of *Sesbania* nitrogen equivalent to 135 kg ha⁻¹ is produced [89].

Green manures which are rich in nitrogen and phosphorus readily decompose, hence were used as a substrate in the form of dry powder. However, individually no single substrate qualifies to be fit for use as soilless media and notably in each there may be limitations. Hence a combination of coirpith (CP), vermicompost (VC), green manure (Daincha) (GM) and sand (S) were used as substrates on the evaluation of growth and yield of ribbed gourd in soilless culture under matric suction irrigation in which crops were raised in growing media in containers that were interconnected by tubs and tubes to maintain moisture always by matric suction (Fig. 10). "Growing media having CP: VC: GM: S with salinity of 0.5–1.0 dS m⁻¹ recorded the highest fruit yield (8.884 kg pot⁻¹) and water use efficiency (312.1 kg fresh fruit mm⁻¹)" [10].



Fig. 10. Effect of salinity and composition of growing media on growth and yield of ribbed gourd in soilless culture under matric suction irrigation. (A) Matric suction assembly (B) Fruiting in growing media CP+VC+GM (1:1:1) [10]

4. CONCLUSION

Legumes including *Sesbania aculeata*, *Sesbana rostrata*, *Vigna radiata* and *Vigna mungo* are the potential green manure crops for their capability of nodule formation and nitrogen fixation. Incorporation of green manures in field remarkably increased soil organic matter, N, P, K and S compared to initial soil status and attained a peak level at 50 days after incorporation. Hence, inclusion of green manures as dry powder in addition to other grow media substrates will be an alternative choice to obtain significant increase in growth and yield parameters of crops in soilless culture. As soilless culture expands research, education, and agricultural opportunities, there are great capabilities to confront these challenges of the green revolution and promote for a soilless one.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. DESA U. World Population Prospects 2022: Summary of Results (UN DESA/POP/2022/TR/NO. 3), United Nations Department of Economic and Social Affairs. Population Division, United Nations, New York. 2022.
2. Leridon H. Population mondiale: vers une explosion ou une implosion?. Population Societes. 2020;573(1):1-4. Available:https://doi.org/10.3917/popsoc.573.0001.
3. Velazquez-Gonzalez RS, Garcia-Garcia AL, Ventura-Zapata E, Barceinas-Sanchez JD, Sosa-Savedra JC. A review on hydroponics and the technologies associated for medium-and small-scale operations. Agriculture. 2022;12(5):646. Available:https://doi.org/10.3390/agriculture12050646.
4. Unicef. The state of food security and nutrition in the world 2018: building climate resilience for food security and nutrition. FAO; 2018.
5. Fedoroff NV. Food in a future of 10 billion. Agriculture & Food Security. 2015;4:1-10. Available:https://doi.org/10.1186/s40066-015-0031-7.
6. Sambo P, Nicoletto C, Giro A, Pii Y, Valentinuzzi F, Mimmo T, Lugli P, Orzes G, Mazzetto F, Astolfi S, Terzano R. Hydroponic solutions for soilless production systems: issues and opportunities in a smart agriculture perspective. Frontiers in plant science. 2019;10:923. Available:https://doi.org/10.3389/fpls.2019.00923.
7. Tzortzakis N, Nicola S, Savvas D, Voogt W. Soilless cultivation through an intensive crop production scheme. Management strategies, challenges and future directions. Frontiers in Plant Science. 2020;11:529970. Available:https://doi.org/10.3389/fpls.2020.00363.
8. Schwarz D, Franken P, Krumbein A, Kläring H, Bar-Yosef B. Nutrient management in soilless culture in the conflict of plant, microorganism, consumer and environmental demands. Acta Hort 2009;843:27-34.
9. Kumar R, Mahajan G, Srivastava S, Sinha A. Green manuring: a boon for sustainable agriculture and pest management—a review. Agricultural Reviews. 2014;35(3):196-206. DOI: 105958/0976-07412014009064
10. Elakiya N, Arulmozhiselvan K. Effect of salinity and composition of growing media on growth and Yield of ribbed gourd in soilless culture under matric suction irrigation. The Pharma Inno. J. 2021;10(10):578-85.
11. Schmilewski G. Growing medium constituents used in the EU. In International Symposium on Growing Media. 2007;2007(819):33-46.
12. Barrett GE, Alexander PD, Robinson JS, Bragg NC. Achieving environmentally sustainable growing media for soilless plant cultivation systems—A review. Scientia horticultrae. 2016;212:220-34.
13. Gruda N, Caron J, Prasad M, Maher MJ. Growing media. In Encyclopedia of Soil

- Sciences. 3rd Edn., R. Lal., ed. 2016;1053–1058.
DOI:<https://doi.org/10.1081/E-ESS3-120053784>.
14. Fussy A, Papenbrock J. An overview of soil and soilless cultivation techniques—chances, challenges and the neglected question of sustainability. *Plants*. 2022;11(9):1153.
 15. Gupta P, Singh J, Verma S, Chandel AS, Bhatla R. Impact of climate change and water quality degradation on food security and agriculture. In *Water Conservation in the Era of Global Climate Change*. Elsevier. 2021;1-22.
 16. Karagöz FP, Dursun A, Karaşal M. A review: use of soilless culture techniques in ornamental plants. *Ornamental Horticulture*. 2022;28:172-80.
 17. Gruda N, Savvas D, Colla G, Roupael Y. Impacts of genetic material and current technologies on product quality of selected greenhouse vegetables—A review. *Eur. J. Hortic. Sci.* 2018;83(11):319-28.
 18. Gruda N, Qaryouti MM, Leonardi C. Growing media. In *good agricultural practices for greenhouse vegetable crops – principles for mediterranean climate areas*. In *Plant Production and Protection Paper (Rome, Italy: FAO)*; 2013.
 19. McGinnis MS, Bilderback TE, Warren SL. Vermicompost amended pine bark provides most plant nutrients for hibiscus moscheutos 'luna blush'. In *International Symposium on Growing Media and Composting 2009*;891:249-256.
 20. Abbey L, Young C, Teitel-Payne R, Howe K. Evaluation of proportions of vermicompost and coir in a medium for container-grown Swiss chard. *International journal of vegetable science*. 2012;18(2):109-20.
 21. Bhat N, Albaho M, Suleiman M, George BT, Ali SI. Growing substrate composition influences growth, productivity and quality of organic vegetables. *Asian J. Agric. Sci.* 2013;5(4):62-6.
 22. Kameswari PL, Girwani A, Padmavathamma AS. Effect of different potting media mixtures on growth and flowering of chrysanthemum (*Dendranthema grandiflora* T.). *Progressive horticulture*. 2014;46(2):314-8.
 23. Yetisir H, Caliskan ME, Soylu S, Sakar M. Some physiological and growth responses of watermelon [*Citrullus lanatus* (Thunb.) Matsum. and Nakai] grafted onto *Lagenaria siceraria* to flooding. *Environmental and experimental botany*. 2006; 58(1-3):1-8.
 24. Majdi Y, Ahmadizadeh M, Ebrahimi R. Effect of different substrates on growth indices and yield of green peppers at hydroponic cultivate. *Current Research Journal of Biological Sciences*. 2012; 4(4):496-9.
 25. Huang L, Gu M, Yu P, Zhou C, Liu X. Biochar and vermicompost amendments affect substrate properties and plant growth of basil and tomato. *Agronomy*. 2020;10(2):224.
 26. Ambayeba Muimba-Kankolongo, Pre- and Postharvest Field Operations, Editor(s): Ambayeba Muimba-Kankolongo, in *Food Crop Production by Smallholder Farmers in Southern Africa*, Academic Press, ISBN 9780128143834. 2018;59-71.
Available:<https://doi.org/10.1016/B978-0-12-814383-4.00007-4>.
 27. Maitra S, Zaman A, Mandal TK, Palai JB. Green manures in agriculture: A review. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(5):1319-27.
 28. Fageria NK, Baligar VC, Bailey BA. Role of cover crops in improving soil and row crop productivity. *Communications in soil science and plant analysis*. 2005;36(19-20):2733-57.
 29. Reddy PP. *Sustainable intensification of crop production*. Singapore: Springer; 2016 Nov 17.
 30. Meena BL, Fagodiya RK, Prajapat K, Dotaniya ML, Kaledhonkar MJ, Sharma PC, Meena RS, Mitran T, Kumar S. Legume green manuring: an option for soil sustainability. *Legumes for soil health and sustainable management*. 2018:387-408.
 31. Delogu G, Cattivelli L, Pecchioni N, De Falcis D, Maggiore T, Stanca AM. Uptake and agronomic efficiency of nitrogen in winter barley and winter wheat. *European Journal of Agronomy*. 1998; 9(1):11-20.
DOI:10.1016/S1161-0301(98)00019-7.

32. Ladha JK, Pareek RP, Becker M. Stem-nodulating legume-Rhizobium symbiosis and its agronomic use in lowland rice. *Advances in Soil Science: Volume 20*. 1992;20:147-92.
DOI:101007/978-1-4612-2930-8_3
33. Sullivan PG, Diver S. Overview of principal uses of cover crops and green manures. *Fundam. Sustain. Agric.* 2003;7:1-6.
34. Becker M, Ladha JK, Ali M. Green manure technology: Potential, usage, and limitations. A case study for lowland rice. In *Management of Biological Nitrogen Fixation for the Development of More Productive and Sustainable Agricultural Systems: Extended versions of papers presented at the Symposium on Biological Nitrogen Fixation for Sustainable Agriculture at the 15th Congress of Soil Science, Acapulco, Mexico*. Springer Netherlands. 1995a;181-194.
35. Adhikary BH, Thakur MK. *Azolla: An alternative source of Chemical Nitrogen in Rice Production*. NARC. 2013.
36. Singh Y, Khind CS, Singh B. Efficient management of leguminous green manures in wetland rice. *Advances in agronomy*. 1991;45:135-89.
37. Kumar S, Samiksha, Sukul P. Green manuring and its role in soil health management. *Soil Health*. 2020:219-41.
38. Becker M, Ali M, Ladha JK and Ottow JCG. Agronomic and economic evaluation of *Sesbania rostrata* green manure establishment in irrigated rice. *Field Crops Res.* 1995b; 40: 135–141.
DOI:10.1016/0378-4290(94)00097-V.
39. Munawar A, Suhartoyo H. Litter production and decomposition rate in the reclaimed mined land under *albizia* and *sesbania* stands and their effects on some soil chemical properties. *Journal of Tropical Soils*. 2013;16(1):1-6.
40. Bouldin DR. In: *Proc. Symposium on Sustainable Agriculture, Int. Rice Res. Inst., Los Banos, Philippines*. 1988;151-163.
41. Meelu OP, Furoc RE, Dizon MA, Morris RA, Marqueses FP. Evaluation of different green manures on rice yield and soil fertility. Paper presented at the 16th annual scientific convention of the Crop Society of Philippines, Central Luzon State University, Munoz, Nueva Ecija, Philippines; 1985.
42. Furoc RE, Dizon MA, Morris RA, Marqueses EP. Effects of flooding regime and planting dates to N accumulation of three *Sesbania* species and consequently to transplanted rice. Paper presented at the 16th annual scientific convention of the crop science Society of Philippines, 8–10 May 1985, Central Luzon State University Munoz, Nueva Ecija, Philippines; 1985.
43. Morris RA, Furoc RE, Dizon MA. Rice Responses to a Short-Duration Green Manure. II. N Recovery and Utilization 1. *Agronomy Journal*. 1986;78(3):413-6.
44. Ladha JK, Watanabe I, Saono S. Nitrogen fixation by leguminous green manure and practices for its enhancement in tropical lowland rice. In: *The International Rice Research Institute (ed) Sustainable agriculture: green manure in rice farming*. IRRI, Los Banos. 1988;165–183.
45. Ghai SK, Rao DL, Lalita B. Comparative study of the potential of *sesbanias* for green manuring. *Tropical Agriculture, UK*. 1985;62(1):52-6.
46. Watanabe I. Use of green manure in northeast Asia. In: *International Rice Research Institute (ed) Organic matter and rice*. IRRI, Los Banos. 1984;229–234.
47. Fried M, Danso S, Zupata F. The methodology of N fixation by non – legumes inferred from field experiments with legumes, *Canadian Journal of Microbiology*. 1983;29:1053-1062.
48. Mann RA, MS Zia and M Salim. New dimensions in green manuring for sustaining the productivity of rice wheat system. *Proceedings of Symposium on Integrated Plant Nutrition Management, NFDC, Islamabad, Pakistan*. Edited by Ahmad N and A. Hamid. 2000;166-185.
49. Saravana PP, Perumal R. Effect of integrated nitrogen management on fertility status of rice soil. *Madras Agricultural Journal*, 2000;87:217-222.
50. Mandal UK, Singh G, Victor US, Sharma KL. Green manuring : its effect on soil properties and crop growth under rice – wheat cropping system, *European Journal of Agronomy*. 2002;19:225-237.

51. Irin IJ, Biswas PK. Residual effect of green manure on soil properties in green manure-transplant aman-mustard cropping pattern. *Indian Journal of Agricultural Research*. 2023;57(1):67-72. DOI:10.18805/IJARE.AF-696.
52. Irin IJ, Biswas PK, Ullah MJ, Roy TS, Khan MA. Influence of green manuring crops on dry matter production and soil health improvement. *Bangladesh Agronomy Journal*. 2019;22(1):39-45.
53. Goyal S, Chander K, Mundra MC, Kapoor KK. Influence of inorganic fertilizers and organic amendments on soil organic matter and soil microbial properties under tropical conditions. *Biology and Fertility of Soils*. 1999;29:196-200. DOI:10.1007/s003740050544.
54. Palaniappan SP. Green manuring: nutrient potentials and management. *Organic Manures and Biofertilizers in Indian Agriculture*. 1991:52-70.
55. Selvi RV, Kalpana R. Potentials of green manure in integrated nutrient management for rice—a review. *Agricultural Reviews*. 2009;30(1):40-7.
56. Sharma AR, Ghosh A. Effect of green manuring with *Sesbania aculeata* and nitrogen fertilization on the performance of direct-seeded flood-prone lowland rice. *Nutrient cycling in Agroecosystems*. 2000;57:141-53. DOI:101023/A:1009863100224.
57. Singh S, Ghoshal N, Singh KP. Synchronizing nitrogen availability through application of organic inputs of varying resource quality in a tropical dryland agroecosystem. *applied soil ecology*. 2007;36(2-3):164-75. DOI:10.1016/j.apsoil.2007.01.007.
58. Sinha AS, Kumar RA, Kamil DE, Kapur PR. Release of nitrogen, phosphorus and potassium from decomposing *Crotalaria juncea* L. in relation to different climatic factors. *Environ Ecol*. 2009;27(4B):2077-81.
59. Yadav RL, Dwivedi BS, Pandey PS. Rice-wheat cropping system: assessment of sustainability under green manuring and chemical fertilizer inputs. *Field crops research*. 2000;65(1):15-30. DOI:101016/S0378-4290(99)00066-0.
60. Von Wandruszka R. Phosphorus retention in calcareous soils and the effect of organic matter on its mobility. *Geochemical transactions*. 2006;7:1-8. DOI: 101186/1467-4866-7-6.
61. Oehl F, Oberson A, Probst M, Fließbach A, Roth HR, Frossard E. Kinetics of microbial phosphorus uptake in cultivated soils. *Biology and Fertility of Soils*. 2001;34:31-41. DOI: 101007/s003740100362.
62. Cavigelli MA, Thien SJ. Phosphorus bioavailability following incorporation of green manure crops. *Soil Science Society of America Journal*. 2003;67(4):1186-94. DOI:10.2136/sssaj2003.1186.
63. Kamh M, Horst WJ, Amer F, Mostafa H, Maier P. Mobilization of soil and fertilizer phosphate by cover crops. *Plant and Soil*. 1999;211:19-27. DOI:101023/A:1004543716488.
64. Sharpley AN, Smith SJ. Mineralization and leaching of phosphorus from soil incubated with surface-applied and incorporated crop residue. *American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America*; 1989. DOI:10.2134/jeq198900472425001800010018x.
65. Tissen H, Cuevas E, Chacon P.. The role of organic matter in sustaining soil fertility *Nature*. 1994;371:783–785
66. Gu C, Lv W, Liao X, Brooks M, Li Y, Yu C, Yang L, Li X, Hu W, Dai J, Zheng W. Green manure amendment increases soil phosphorus bioavailability and peanut absorption of phosphorus in red soil of South China. *Agronomy*. 2023;13(2):376. Available:https://doi.org/10.3390/agronomy13020376.
67. Shen J, Li H, Neumann G, Zhang F. Excess cation uptake, and extrusion of protons and organic acid anions by *Lupinus albus* under phosphorus deficiency *Plant Science*. 2005;168:837-845 DOI:101016/S0168-9452(01)00373-9.
68. Sas L, Rengel Z, Tang C. Excess cation uptake, and extrusion of protons and organic acid anions by *Lupinus albus*

- under phosphorus deficiency. *Plant Science*. 2001;160(6):1191-8.
DOI:10.1016/S0168-9452(01)00373-9.
69. Islam MM, Urmi TA, Rana MS, Alam MS, Haque MM. Green manuring effects on crop morpho-physiological characters, rice yield and soil properties. *Physiology and Molecular Biology of Plants*. 2019; 25:303-12.
 70. Rani TS, Ramulu C, Kumar TS, Rao PJ. Evaluation of Dhaincha (*Sesbania aculeata* L.) accessions for green manuring traits and soil fertility improvement. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(6):1932-6.
 71. Talgre L, Luringson E, Roostalu H, Astover A, Makke A. Green manure as a nutrient source for succeeding crops; 2012.
 72. Bhuiyan NI, Zaman SK. *Biological Nitrogen Fixation Associated with Rice Production*. Kluwer Academic Publishers, The Netherlands; 1996.
 73. Becker M, Ladha JK, Orrow JCG. Growth and N₂ Fixation of Two Stem-Nodulating Legumes and Their Effect as Green Manure on Lowland Rice. *Soil Biol and Biochem*. 1992;22:1109–1119.
DOI:10.1016/0038-0717(90)90037-Z
 74. Bhattacharya K, Mandal SR. Effect of green manuring on availability and losses of plant nutrients by soil. *Indian Agriculturist*. 1996;40:199-208.
 75. Eriksen J, Thorup-Kristensen K. The effect of catch crops on sulphate leaching and availability of S in the succeeding crop on sandy loam soil in Denmark. *Agri Eco and Env*. 2002;90:247–254.
DOI:10.1016/S0167-8809(01)00214-6
 76. Rumball W. *Grasslands Puna'chicory (Cichorium intybus L.)*. New Zealand journal of experimental agriculture. 1986; 14(1), 105-107.
 77. MacRae RJ, Mehuys GR. The effect of green manuring on the physical properties of temperate area soils. *Adv in Soil Sci*. 1985;3:71–94.
DOI:10.101007/978-1-4612-5090-6_2.
 78. Carvalho NS, Oliveira AN, Calaã MM, Neto VP, de Sousa RS, dos Santos VM, de Araujo AS. Short-term effect of different green manure on soil chemical and biological properties. *African Journal of Agricultural Research*. 2015;10(43):4076-81.
 79. Bhuiyan NI, Zaman SK, GM Panaullah. Dhaincha green manure: a potential nitrogen source for rainfed lowland rice. *Proceedings of the Workshop on Experiences with Modern Rice Cultivation in Bangladesh*; 1988.
 80. Yang L, Bai J, Liu J, Zeng N, Cao W. Green Manuring Effect on Changes of Soil Nitrogen Fractions, Maize Growth, and Nutrient Uptake. *Agronomy*. 2018;8(11):261.
Available:https://doi.org/10.3390/agronomy8110261.
 81. Rahman MH, Islam MR, Jahiruddin M, MQ Haque. Management of organic manure and inorganic fertilizer in the maize-mungbean/dhaincha-T. aman rice cropping pattern for increased crop production. *Bangladesh Journal of Agricultural Research*. 2012;37(2):225-234.
 82. Mann RA, Zia MS, Saleem M. An improved green manure technology for sustaining the wheat rice system. *Quarterly Sci. Vision*. 2000;6(2):53.
 83. Sarwar AK. Characterization of dhaincha accessions based on morphological descriptors and biomass production. *Journal of the Bangladesh Agricultural University*. 2015;13(1):49-54.
 84. Rahman MH, Islam MR, Jahiruddin M, Rafii MY, Hanafi MM, Malek MA. Integrated nutrient management in maize-legume-rice cropping pattern and its impact on soil fertility. *J. Food Agric. Environ*. 2013;11(2):648-52.
 85. Ankur Singh, Mohan Lal, Shivashankar K, Kunwar Satyendra Singh, Richa Tiwari. Effect of natural farming practices on growth and yield of black gram under pigeonpea + black gram cropping system. *The Pharma Innovation Journal* 2023;12(3):5871-5874
DOI:10.13140/RG.2.2.17790.46408.
 86. Ugale NS. Effect of different organic inputs with jeevamrut on yield, quality and soil properties in soybean-wheat cropping sequence (Doctoral dissertation, PhD Dissertation, Mahatma Phule Krishi Vidyapeeth, Rahuri, India); 2014.

87. Kumbar BA, Devakumar N. Effect of jeevamrutha and Panchgavya on growth, yield and microbial population of french bean (*Phaseolus vulgaris* L.). *Advanced Life Sciences*. 2016; 5:3619-23.
88. Sutar R, Sujith GM, Devakumar N. Growth and yield of Cowpea [*Vigna unguiculata* (L.) Walp] as influenced by jeevamrutha and panchagavya application. *Legume Research-An International Journal*. 2019; 2(6):824-828.
89. Ehsan S, Niaz A, Saleem I, Mehmood K. Substitution of major nutrient requirement of rice-wheat cropping system through *Sesbania* green manuring. *Sci. Agric*. 2014;8:99-102.
90. Noor-A-Jannat S Ahmed, Abuyusuf M, Hassan MZ, Lipi NJ, Biswas KK. Nitrogen fertilizer after green manuring on the yield of t. aman rice. *American Research Thoughts* 2015;1:2898-2909.
91. Geethalakshmi V. Studies on the direct and residual effect of non-conventional green leaf manures with n for rice-rice cropping system. Tamil Nadu Agricultural University; Coimbatore; 1996.
92. Sarwar AK, Hossain SZ, Chanda SC. Effect of Dhaincha accessions on soil health and grain yield of rice. *Journal of Bioscience and Agriculture Research*. 2017;13(02):1140-5.
93. Ajaykumar R, Venkitaswamy R, Rajeshkumar A. "Sustainable Nutrient Management Practices to Increase the Productivity of Transplanted Lowland Rice." *Madras Agricultural Journal*. 2017;104(7/9):226-229.
94. Chanda SC, Rafiqul Islam M, Golam Sarwar AK. Organic Matter Decomposition and Nutrient Release from Different Dhaincha (*Sesbania* spp.) Genotypes. *Journal of Agricultural Sciences (Sri Lanka)*. 2021;16(2).
95. Ahmed P, Nath RK, Sarma R. Cultivation of green manuring crops for improving soil health and increasing yield of rice in Tinsukia district of Assam-a case study. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(2):655-7.
96. Thorup-Kristensen K, Magid J, Jensen LS. Catch crops and green manures as biological tools in nitrogen management in temperate zones. *Adv. Agron*. 2003;79:227–302.
97. Mandal UK, Singh G, Victor US, Sharma KL. Green manuring: its effect on soil properties and crop growth under rice–wheat cropping system. *European Journal of Agronomy*. 2003;19(2):225-37.
98. Phiri E, Verplancke H, Kwesiga F, Mafongoya P. Water balance and maize yield following improved *sesbania* fallow in eastern Zambia. *Agroforestry systems*. 2003;59:197-205.
99. Adekiya AO, Agbede TM, Aboyeji CM, Dunsin O, Ugbe JO. Green manures and NPK fertilizer effects on soil properties, growth, yield, mineral and vitamin C composition of okra (*Abelmoschus esculentus* (L.) Moench). *Journal of the Saudi Society of Agricultural Sciences*. 2019;18(2):218-23.
100. Lee CR, Kim SH, Oh Y, Kim YJ, Lee SM. Effect of Green Manure on Water-Stable Soil Aggregates and Carbon Storage in Paddy Soil. *Korean Journal of Soil Science and Fertilizer*. 2023;56(2):191-8.
101. Dhar D, Datta A, Basak N, Paul N, Badole S, Thomas T. Residual effect of crop residues on growth, yield attributes and soil properties of wheat under rice-wheat cropping system. *Indian Journal of Agricultural Research*. 2014;48(5):373-8.
102. Adekiya AO, Agbede TM, Aboyeji CM, Dunsin O, Ugbe JO. Green manures and NPK fertilizer effects on soil properties, growth, yield, mineral and vitamin C composition of okra (*Abelmoschus esculentus* (L.) Moench). *Journal of the Saudi Society of Agricultural Sciences*. 2019 Apr 1;18(2):218-23.
Available:<https://doi.org/10.1016/j.jssas.2017.05.005>.
103. Hundal HS, Biswas CR, Vig AC. Phosphorus sorption characteristics of flooded soil amended with manures. *Tropical agriculture*.
104. Ye X, Liu H, Li Z, Wang Y, Wang Y, Wang H, Liu G. Effects of green manure continuous application on soil microbial biomass and enzyme activity. *Journal of Plant Nutrition*. 2014;37(4):498-508.
105. Eriksen J. Gross sulphur mineralisation–immobilisation turnover in soil amended with plant residues. *Soil biology and Biochemistry*. 2005;37(12):2216-24.
DOI:10.1016/j.soilbio.2005.04.003.

106. Sikora LJ, Stott DE. Soil organic carbon and nitrogen Soil Science Society of America. 1996.
107. Doran JW, Fraser DG, Culik MN, Liebhardt WC. Influence of alternative and conventional agricultural management on soil microbial processes and nitrogen availability. *American Journal of alternative agriculture*. 1987;2(3):99-106.
DOI:10.1017/S0889189300001739.
108. Kautz T, Wirth S, Ellmer F. Microbial activity in a sandy arable soil is governed by the fertilization regime. *European Journal of Soil Biology*. 2004;40(2):87-94.
DOI:10.1016/j.ejsobi.2004.10.001
109. Campbell CA, Biederbeck VO, Zentner RP, Lafond GP. Effect of crop rotations and cultural practices on soil organic matter, microbial biomass and respiration in a thin Black Chernozem. *Canadian Journal of Soil Science*. 1991;71(3):363-76.
110. Schutter M, Dick R. Shifts in substrate utilization potential and structure of soil microbial communities in response to carbon substrates. *Soil Biology and Biochemistry*. 2001 Sep 1;33(11):1481-91.
DOI:10.1016/S0038-0717(01)00057-8.
111. Liu GS, Luo ZB, Wang Y, Li HL, Wang GF, Ma JM. Effect of green manure application on soil properties and soil microbial biomass in tobacco field. *Journal of Soil Water Conservation*. 2006;20:95 – 98.
112. Hu C, Cao ZP, Ye ZN, Wu WL. Impact of soil fertility maintaining practice on soil microbial biomass carbon in low production agro-ecosystem in northern China. *Acta Ecologica Sinica*. 2006;26(3):808-14.
113. Swarup, A. "Effect of *Sesbania bispinosa* decomposition time and sodality on rice yield." *AGRIS Agricultural Information Management Standards*. 1991;13:28-29.
114. Odhiambo JJ. Decomposition and nitrogen release by green manure legume residues in different soil types. *African Journal of Agricultural Research*. 2010;5(1):090-6.
115. Chanda Sontosh C, Golam Sarwar AKM. Status of dhaincha incorporated soil after rice harvest in (boro) rice–dhaincha–rice (t. aman) cropping pattern. 2017.
116. Shi J. Decomposition and nutrient release of different cover crops in organic farm systems; 2013.
117. Selvi RV, Nadanassababady T, Rajendran P. Green manuring in lowland rice—a review. *Agricultural Reviews*. 2005;26(1):14-26.
118. Th o" nnissen C, Midmore DJ, Ladha JK, Olk DC, Schmidhalter U. Legume decomposition and nitrogen release when applied as green manures to tropical vegetable production systems. *Agronomy Journal*. 2000;92(2):253-60.
119. Xu J, Si L, Zhang X, Cao K, Wang J. Various green manure-fertilizer combinations affect the soil microbial community and function in immature red soil. *Frontiers in Microbiology*. 2023;14:1255056.
DOI:10.3389/fmicb.2023.1255056.
120. Clement A, Ladha JK, Chalifour FP. Nitrogen dynamics of various green manure species and the relationship to lowland rice production. *Agronomy journal*. 1998;90(2):149-55.
121. Dux J, Norgrove L, Hauser S, Wick B, Kuhne R. Plant leaf residue decomposition, nutrient release and soil enzyme activity. Conference on International Agricultural Research for Development: Prosperity and poverty in a globalised world—Challenges for agricultural research; 2006.
122. Berg B, Steffen KT, McClaugherty C. Litter decomposition rate is dependent on litter Mn concentrations. *Biogeochemistry*. 2007;82:29-39.
123. Setiawati MR, Prayoga MK, Stöber S, Adinata K, Simarmata T. Performance of rice paddy varieties under various organic soil fertility strategies. *Open agriculture*. 2020;5(1):509-15.
124. Kabir AK, Moniruzzaman M, Gulshan Z, Rahman AB, Sarwar AK. Biomass Yield, Chemical Composition and In Vitro Gas Production of Different Dhaincha (*Sesbania* spp.) Accessions from Bangladesh. *Indian Journal of Animal Nutrition*. 2018;35(4):397-402.
125. Liu X, Xie Z, Wang Y, Sun Y, Dang X, Sun H. A dual role of amino acids from *Sesbania rostrata* seed exudates in the

- chemotaxis response of Azorhizobium caulinodans ORS571. *Molecular Plant-Microbe Interactions*. 2019;32(9):1134-47.
126. Chanda S, Hossain M, Uddin M, Islam M, Sarwar AG. Fiber Yield, Physical and Biochemical Properties of Three Species of Sesbania. *Bangladesh Agron. J.* 2019;21:79-85.
127. Hossain MA, Becker K. Nutritive value and antinutritional factors in different varieties of Sesbania seeds and their morphological fractions. *Food Chemistry*. 2001;73(4):421-31.
128. Xie Z, Tu S, Shah F, Xu C, Chen J, Han D, Liu G, Li H, Muhammad I, Cao W. Substitution of fertilizer-N by green manure improves the sustainability of yield in double-rice cropping system in south China. *Field Crops Research*. 2016;188:142-9.
129. Palaniappan SP, Siddeswaran K. Regional overview on green manure in rice-based cropping systems. Legumes in rice-based cropping systems in tropical Asia: constraints and opportunities. 2001; 126-135.
130. Chavan PD, Karadge BA. Growth, mineral nutrition, organic constituents and rate of photosynthesis in Sesbania grandiflora L. grown under saline conditions. *Plant and Soil*. 1986;93:395-404.
131. Carroll Andrew, Chris Somerville. "Cellulosic biofuels." *Annual review of plant biology*. 2009;60:165-182.
132. Factsheet. *Tropical Forages. Sesbania sesban*. 2016;1-7.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/118987>