



Assessment of Bees' Diversity within Ngel-Nyaki Forest Reserve at Taraba State, Nigeria

L. C. Garba ^a, D. L. David ^a, E. Elisha ^b, S. D. Titus ^a,
J. H. Bulus ^a and J. P. Yani ^{c*}

^a Department of Biological Sciences, Taraba State University, Jalingo, Taraba State, Nigeria.

^b Nigerian Montane Forest Project, Ngel-Nyaki, Taraba State, Nigeria.

^c Department of Forestry and Wildlife, Taraba State University, Jalingo, Taraba State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRAF/2023/v9i3213

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/99643>

Review Article

Received: 04/04/2023

Accepted: 27/05/2023

Published: 17/06/2023

ABSTRACT

A field survey was conducted in Ngel-Nyaki Forest Reserve to assess the diversity of bees within the Reserve. Two methods of sampling techniques were used to sample bee species, opportunistic sampling and systematic sweep netting. Opportunistic sampling method was applied by sweep netting bees along 100 m plot where flowering plants were blooming. Collection was conducted by a single collector using an active search by walking randomly around the plot observing flowering plants and searching for bees foraging on the flowers then a sweep net was used for capturing bees. Sweep netting was conducted, line transect established within the plots for two hours, from 6:00am to 11:00am spending 20 minutes in each plot considering when the weather is conducive for bees' activity. This was done using entomological sweep net. Bees captured were killed in a killing bottle containing 70% ethanol. Specimens were pinned on a board and unique codes were assigned to each species before identification. Bees were identified to species level by using

*Corresponding author: E-mail: yanij0163@gmail.com;

published systematic keys, Bees field guide and use of online materials. All data were computed using Microsoft® Excel® 2016 and percentage abundance composition of bees were calculated. 575 bees spreading across 3 families and 12 species were sampled. The most abundant species was the honey bee (*Apis mellifera*) (245; 42.53%), followed by mimic bee (*Megachile* sp.) (89; 15.45%), and the least was leafcutter bee (*Osmia* sp.) (2; 0.35%). In terms of diversity by species, Shannon Diversity Index (H) and Shannon Equitability Index (EH) were 1.694893 and 0.736083 respectively while diversity by plots using Shannon Diversity Index (H) and Shannon Equitability Index (EH) were 1.750688 and 0.977078 respectively. This study therefore, brings to the fore the diversity of bees within Ngel-Nyaki Forest Reserve and underline the need for sustainable actions to be taken to conserve beneficial rare species while managing the abundant ones.

Keywords: Assessment; diversity; forest; Ngel-Nyaki.

1. INTRODUCTION

Bees belong to the Order Hymenoptera, Sub-order Apocrita, Super-family Apoidea, Clade Anthophila and are of monophyletic lineage [1,2]. There are about 20,900 known species of bees in eleven (11) recognized biological families [3], though many are yet to be identified. Out of the eleven (11) bee families, six (6) are found in Africa [4] namely, Apidae, Megachilidae, Melitidae, Andrenidae, Colletidae and Halictidae [5,6]. They are found on every continent except Antarctica, in every habitat on the planet that contains insect pollinated flowering plants [7] (Masigà, 2017). Bees range in size from tiny stingless bee species (*Trigonaminima*) (Hymenoptera, Stenotritidae), whose workers are less than 2 millimeters long [8], to large species of leafcutter bee (*Megachilepluto*) (Hymenoptera, Melitidae) [9], whose females can attain a length of 39 millimeters [10,11]. Bees are among the main pollinators in most ecosystems, as they rely on flowers for proteins [12,13], oils and sugar throughout their life cycle [10]. They also exhibit the highest floral visitation rates among all the insects (Masigà, 2017). There is little study on the different species of bees in Nigeria [14] as most of the work conducted in Nigeria has only been on honey bees and bee keeping. Honey bees are just one out of many species of bees [15]. So also, the Ngel-Nyaki Forest Reserve is a biodiversity hotspot zone with different species of bees but till date there is no documented work on the taxonomy and different species of bees found within the Reserve [16]. The findings of the study therefore, will contribute in the field of conservation, by identifying the different species of bees within Ngel-Nyaki Forest Reserve. The work will also help to document the different types of bees within the Forest Reserve by creating a checklist for the different species of bees as well as the percentage abundance

composition of bees found within Ngel-Nyaki Forest Reserve.

1.1 Study Area

Ngel-Nyaki is located towards the western escarpment of the Mambilla Plateau. The Forest Reserve forms part of the Guinean forest biodiversity hotspot which extends from Sierra Leone to Cameroon Mountains. The reserve is situated between longitude 07° 05'N and latitudes 011° 05'E at an altitude of 1,450m–1,600m above sea level (Fig. 1). The Reserve occupies about 46km² of land, montane grassland, forest fragments, woodland forests and lowlands with about 7.2km² of submontane to mid-altitude forest [17,18]. The Forest Reserve can be reached on foot 40 minutes from Yelwa village [19]. Ngel-Nyaki Forest Reserve is one of the most diverse sub-montane forests in Nigeria. It contains many plants which are endemic to the Afro-montane region. There are more than 146 vascular plants out of which 75 are in IUCN Red List [18]; 4 endangered species and 8 new species. The forest is an Important Bird Area (IBA) [20] and has been classified as an Important Bird Area by Bird Life International and a critical site for biodiversity conservation [21]. The forests fall within the Cameroon Highlands Eco region, which are exceptionally rich in endemic taxa. Ngel-Nyaki forest, a 46 km² forest reserve, is the most species rich forest on the Plateau and is the location of the Nigerian Montane Forest Project field station. Ngel-Nyaki forest harbours 9 IUCN Red Data Listed tree species with at least another 20 other species in the genera *Warneckia* and *Deinbollia*, new to science. It is rich in primate species, including the IUCN Endangered/Nigeria-Cameroon chimpanzee *Pan troglodyteselliotti*. There are African golden cats *Caracal aurata*, 3 new species of frogs that the Nigerian Montane

Forest Project have identified so far. However, Ngel-Nyaki and other Mambilla Plateau forests and the animal species are threatened because of increased human and cattle populations, poverty and climate change. Fulani pastoralists are increasingly moving their cattle into the forests and burning forest for grazing, farmers are felling trees and planting crops and poachers are hunting wildlife [19].

2. MATERIALS AND METHODS

2.1 Sampling Methods

Six (6) plots were established randomly where flowering plants were found within the Forest Reserve after a pilot survey.

2.1.1 Bee sampling

Two (2) methods of sampling techniques were used to sample bee species. These were Opportunistic sampling and Systematic sweep netting. Opportunistic sampling method was

applied by sweep netting bees along 100 m plot where flowering plants were blooming. Collection was conducted by a single collector using an active search by walking randomly around the plot observing flowering plants and searching for bees foraging on the flowers then a sweep net was used for capturing bees [22]. Sweep netting was conducted, line transect established within the plots for 2 hours, from 6:00am to 11:00am spending 20 minutes in each plot considering when the weather is conducive for bees' activity. This was done using entomological sweep net. Bees captured from flowers were killed in a killing bottle containing 70% ethanol. Specimens were pinned on a board and unique codes were assigned to each species before identification [23-26].

2.2 Identification

Bees were identified to species level using published systematic keys, Bee's field guide and use of online materials [27-29].




Fig. 1. Map of Ngel-Nyaki Forest Reserve, Taraba State, Nigeria
[Source: Nigerian Montane Forest Project (NMFP)].

2.3 Data Analysis

All data were computed using Microsoft® Excel® 2016 and percentage abundance composition of bees were calculated using the formula below by Potts et al. [30] and LeFeon et al. [31]:

$$\% \text{ Abundance COmposition} = \frac{ns}{Ns} \times 100$$

Where ,

'ns' is the number of individual species.

'Ns' is the total number of species captured during the survey.

3. RESULTS AND DISCUSSION

At the end of the survey, 575 bees were captured comprising 12 species namely; Leafcutter Bee (*Ammobatodes scriptus*), Mining Bee (*Andrena cineraria*), Honey Bee (*Apis mellifera*), Bumble Bee (*Bombus sp.*), Memic Bee (*Megachile spp.*), Stingless Bee (*Meliponulabeccarii*), Manson Bee (*Osmia caerulea*), Leafcutter Bee (*Osmia sp.*), Carpenter Bee (*Xylocopa augusti*), Carpenter Bee (*Xylocopa caffra*), Violet Bee (*Xylocopa violacea*) and Squash Bee (*Penonapis sp.*). They were captured in the six (6) plots within the Reserve and identified. The most abundant species was the Honey Bee (*Apis*

mellifera) with 245 individuals followed by Memic Bee with 89 individuals then Stingless Bee with 70 individuals while the least was *Osmia sp.* with 2 individuals captured during the survey. Plot 3 (144) captured the highest number of bees, followed by Plot 1 (112), then Plot 2 (104) and the least was Plot 4 recording 65 bees.

In terms of diversity by species, Shannon Diversity Index (H) and Shannon Equitability Index (EH) were 1.694893 and 0.736083 respectively while diversity by plots using Shannon Diversity Index (H) was 1.750688 and Shannon Equitability Index (EH) was 0.977078.

This information is shown on Table 1.

Table 2 represents the number of bee species according to their families. All the bees sampled were members of three (3) families' namely; Andrenidae, Apidae and Megachilidae. Apidae (443) took the lead, followed by Megachilidae (120) and the least was Andrenidae (13).

The percentage composition of bees varied across the three (3) families and genera. Honey Bee recorded the highest percentage composition of 42.53%, followed by Memic Bee (15.45%), then Stingless Bee (12.15%) and the least was Leafcutter Bee (*Osmia sp.*) (0.35%) as shown below in Table 3.

Table 1. Number of bees captured in six (6) plots during the survey

S/ N	Common Name	Scientific name	Plot 1	Plot 2	Plot 3	Plot 4	Plot 5	Plot 6	Total
1	Leaf cutter Bee	<i>Ammobatodes scriptus</i>	3	-	5	5	-	-	13
2	Mining Bee	<i>Andrena cineraria</i>	6	4	-	3	-	-	13
3	Honey Bee	<i>Apis mellifera</i>	40	70	65	15	15	40	245
4	Bumble Bee	<i>Bombus sp.</i>	7	-	13	9	-	3	32
5	Memic Bee	<i>Megachile sp.</i>	9	10	30	19	13	8	89
6	Stingless Bee	<i>Meliponulabeccarii</i>	16	-	4	-	45	5	70
7	Manson Bee	<i>Osmia caerulea</i>	-	-	12	4	-	-	16
8	Leafcutter Bee	<i>Osmia sp.</i>	2	-	-	-	-	0	2
9	Carpenter Bee	<i>Xylocopa augusti</i>	-	9	5	-	11	-	25
10	Carpenter Bee	<i>Xylocopa caffra</i>	7	-	10	-	-	10	27
11	Violet Bee	<i>Xylocopa violacea</i>	19	11	-	10	-	-	40
12	Squash Bee	<i>Penonapis sp.</i>	3	-	-	-	-	-	3
Total			112	104	144	65	84	66	575

Diversity by species: Shannon Diversity Index (H) = 1.694893, Shannon Equitability Index (EH) = 0.736083; Diversity by plots: Shannon Diversity Index (H) = 1.750688, Shannon Equitability Index (EH) = 0.977078

Table 2. Number of bee species according to family

S/N	Family Name	Common Name	Scientific Name	Number
1	Andrenidae	Mining Bee	<i>Andrenidae cineraria</i>	13
2	Apidae	Bumble Bee	<i>Bombus sp.</i>	32
		Honey Bee	<i>Apis mellifera</i>	245
		Stingless Bee	<i>Meliponulabeccarii</i>	70
		Carpenter Bee	<i>Xylocopa augusti</i>	25
		Carpenter Bee	<i>Xylocopa caffra</i>	27
		Violet Bee	<i>Xylocopa violacea</i>	41
		Squash Bee	<i>Penonapiss.p</i>	3
		Total		443
3	Megachilidae	Leafcutter Bee	<i>Osmia sp.</i>	2
		Manson Bee	<i>Osmia caerulescens</i>	16
		Leafcutter Bee	<i>Ammobatodes scriptus</i>	13
		Memic Bee	<i>Megachile sp.</i>	89
		Total		120

Table 3. Percentage composition of bees' species in the study area

Common name	Scientific name	Percentage composition (%)
Leafcutter Bee	<i>Ammobatodes scriptus</i>	2.00
Mining Bee	<i>Andrena cineraria</i>	2.00
Honey Bee	<i>Apis mellifera</i>	42.53
Bumble bee	<i>Bombus sp.</i>	5.56
Memic Bee	<i>Megachile sp.</i>	15.45
Stingless Bee	<i>Meliponulabeccarii</i>	12.15
Manson Bee	<i>Osmia caerulescens</i>	2.78
Leafcutter Bee	<i>Osmia sp.</i>	0.35
Carpenter Bee	<i>Xylocopa augusti</i>	4.34
Carpenter Bee	<i>Xylocopa caffra</i>	4.69
Violet Bee	<i>Xylocopa violacea</i>	7.12
Squash Bee	<i>Penonapissp</i>	0.52
Total		100%

3.1 Discussion

Brosi et al. [24] and McGavin [23] reported that targeted sweep netting was the best method of sampling bees. This assertion is in agreement with the present study which recorded large populations of bees' species collected using the same method. In the current study, only three (3) families of bees were found within the Reserve, this is in contrast to the six (6) families previously reported by Michener [10] and Masigà (2017) that out of the eleven (11) bees' families, six (6) are found in Africa. The carpenter bees (*Xylocopa sp.*) were more diversified. Bees rely on flowers for protein, oils and sugar throughout their life cycle [10]. All bees in the study area were captured on flowering plants. It was also observed during the survey that as flowering plants increased so also the number of bees increased, this agrees with the report by Kasina et al. [32] who reported that bees need plants for other uses such as getting nest materials, hiding, mating or just as resting sites.

Out of the six (6) plots created within the Reserve, plot three (3) had most of the bees' species as almost all the species were found in the plot, this was because all plants in the plot were in bloom during the survey, while plot 4 had least species because the plot was on high elevation; this agrees with the report by Lefebvre et al. [33] who reported that abundance and diversity of pollinators decrease with increasing elevation. The honey bees (*Apis mellifera*) were found across all the six (6) plots making them more abundant than the other species.

Percentage abundance composition also showed that honey bee (*Apis mellifera*) is the most abundant species with about 245 (42.53%) individuals captured; this is because honey bees are social insects and live in colony with each worker visiting flowers 8-10 times daily in search of pollens [34] and the least was *Osmia sp.* with only 2(0.31%) individuals captured during the survey, probably because it is uncommon and rare.

4. CONCLUSION

In conclusion, the present survey has shown that Ngel-Nyaki Forest Reserve is rich in bees' diversity. It has also documented probably for the very first time, the bees' fauna in the Forest Reserve. At the end of the survey a total of 12 different bee species were caught belonging to three (3) bees' family. Result obtained showed that honey bee (*Apis mellifera*) was the most abundant bee species found within the Reserve while the least was the *Osmia sp.*

ACKNOWLEDGEMENT

We want to first of all acknowledge TETFund's Sponsorship through the Management of Taraba State University, Jalingo for awarding us the Grant and the Management of the Nigerian Montane Forest Project (NMFP) Ngel-Nyaki, Nigeria for their field assistance, materials and accommodation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kayaalp P. The Diversification, biogeography and body size evolution of Australian hylaeine and Euryglossine bees. Flinders University, School of Biological Sciences; 2011.
2. Masiga R, Kasina JM, Mbugi J, Odhiambo C, Kinuthia W, Gemmill-Herren B, Vaissière BE. Do French beans (*Phaseolus vulgaris* L.) grown in proximity to Mount Kenya forest in Kenya experience pollination deficit? *Journal of Pollination Ecology*. 2014;14(24):255-260.
3. Ojwang MRD. Diversity and abundance of bees and their effect on yield of french beans adjacent Mt Kenya forest in Laikipia County. Unpublished Doctoral dissertation, Kenyatta University; 2017.
4. Eardley C, Kuhlmann M, Pauly A. The bee genera and subgenera of sub-Saharan Africa. Brussels: Belgian Development Cooperation. 2010:145.
5. Buchmann SL. The ecology of oil flowers and their bees. *Annual Review of Ecology and Systematics*. 1987;18(1):343-369.
6. Muli E, Patch H, Frazier M, Frazier J, Torto B, Baumgarten T, Grozinger C. Evaluation of the distribution and impact of parasites, pathogens and pesticides on honey bee (*Apis mellifera*) populations in East Africa. *Plos One*. 2014;9(4):e94459.
7. Gilgert W, Vaughan M. The value of pollinators and pollinator habitat to rangelands: Connections among pollinators, insects, plant communities, fish, and wildlife. *Rangelands*. 2011;33(3):14-19.
8. Asiko GA. Pollination of strawberry in Kenya by stingless bees (Hymenoptera: Meliponini) and honey bee (Hymenoptera: Apini) for improved fruit quality. Unpublished Doctoral dissertation, University of Nairobi, Kenya; 2012.
9. Willmer PG, Stone GN. Behavioral, ecological and physiological determinants of the activity patterns of bees. *Advances in the Study of Behavior*. 2004;34(34):347-466.
10. Michener CD. The bees of the world. First edition, Johns Hopkins University Press, Baltimore, M. D. 2000:928.
11. Ascher J, Eardley GC, Mello PT, Ruggiero A, Williams M, Walter P, Warrit K. Integrated taxonomic information system; 2008. Available:<http://www.government/beechecklist.html>
12. Roulston TAH, Cane JH, Buchmann SL. What governs protein content of pollen: pollinator preferences, pollen–pistil interactions, or phylogeny? *Ecological Monographs*. 2000;70(4):617-643.
13. Russo L, Vaudo AD, Fisher CJ, Grozinger CM, Shea K. Bee community preference for an invasive thistle associated with higher pollen protein content. *Oecologia*. 2019;190(4):901-912.
14. Abdullahi G, Sule H, Chimoya IA, Isah MD. Diversity and relative distribution of honeybees foraging plants in some selected reserves in Mubi region, Sudan savannah ecological zone of Nigeria. *Advances in Applied Science Research*. 2011;2(5):388-395.
15. Nsor CA. Sunbird pollination and the fate of strong contributors to a mutualistic network in a West African Montane Forest; 2014.
16. Cheek M, Onana JM, Chapman HM. The montane trees of the Cameroon Highlands, West-Central Africa, with *Deinbolliaonanae* sp. nov. (Sapindaceae), a new primate-dispersed, endangered species. *Peer Journal*. 2021;9:e11036.

17. Korndoerfer MC. Exploring human-environment interactions and their effects around Ngel-Nyaki Forest Reserve, Nigeria; 2010.
18. Borokini TI, Babalola FD, Amusa TO, Ivande ST, Wala ZJ, Jegede OO, Ihuma JO. Tropical montane forest biodiversity in Nigeria—case study of Ngel-Nyaki Forest Reserve, Mambilla Plateau. *International Journal of Environmental Science*. 2012;1(2):95-104.
19. Chapman JD, Chapman HM. The forests of Taraba and Adamawa states, Nigeria: An ecological account and plant species checklist. Department of Plant and Microbial Sciences, University of Canterbury, New Zealand. 2001:146.
20. Ihuma J, Chiuma UD, Chapman HM. Tree species diversity in a Nigeria Montane Forest ecosystem and adjacent fragmented forests. *ARPN Journal of Agricultural and Biological Science* 2010;6(2):17-22.
21. Ezealor AU. Critical sites for biodiversity conservation in Nigeria. Nigerian Conservation Foundation, Lagos. 2002:110.
22. Nielsen A, Steffan-Dewenter I, Westphal C, Messinger O, Potts SG, Roberts SP, Petanidou T. Assessing bee species richness in two Mediterranean communities: importance of habitat type and sampling techniques. *Ecological Research*. 2011;26(5):969-983.
23. McGavin GC. Expedition field techniques: Insects and other terrestrial arthropods. London: Geography Outdoors: The Center Supporting Field Research, Exploration and Outdoor Learning, 1997:1–96.
24. Brosi BJ, Daily GC, Shih TM, Oviedo F, Durán G. The effects of forest fragmentation on bee communities in tropical countryside. *Journal of Applied Ecology*. 2008;45(3):773–783.
25. Westphal C, Bommarco R, Carré G, Lamborn E, Morison N, Petanidou T, Steffan-Dewenter I. Measuring bee diversity in different European habitats and biogeographical regions. *Ecological Monographs*. 2008;78(4):653-671.
26. Prendergast KS, Menz MH, Dixon KW, Bateman PW. The relative performance of sampling methods for native bees: an empirical test and review of the literature. *Ecosphere*. 2020;11(5):e03076.
27. Francoy TM, Silva RAO, Nunes-Silva P, Menezes C, Imperatriz-Fonseca VL. Gender identification of five genera of stingless bees (Apidae, Meliponini) based on wing morphology. *Genetics and Molecular Research*. 2009;8(1):207-214.
28. Buschbacher K, Ahrens D, Espeland M, Steinhage V. Image-based species identification of wild bees using convolutional neural networks. *Ecological Informatics*. 2020;55:101017.
29. Trianto M, Purwanto H. Morphological characteristics and morphometrics of stingless bees (Hymenoptera: Meliponini) in Yogyakarta, Indonesia. *Biodiversitas Journal of Biological Diversity*. 2020;21(6).
30. Le Féon V, Burel F, Chifflet R, Henry M, Ricroch A, Vaissiere BE, Baudry J. Solitary bee abundance and species richness in dynamic agricultural landscapes. *Agriculture, Ecosystems and Environment*. 2013;166:94-101.
31. Potts SG, Vulliamy B, Dafni A, Ne'eman G, Willmer P. Linking bees and flowers: How do floral communities structure pollinator communities? *Ecology*. 2003;84(10):2628-2642.
32. Kasina JM. Bee pollinators and economic importance of pollination in crop production: Case of Kakamega, Western Kenya. Ecology and Development Series No. 54. Dissertation, University of Bonn; 2007.
33. Lefebvre V, Vilemant C, Daugeron C. Altitudinal, temporal and trophic partitioning of flower-visitors in Alpine communities. *Scientific Report*. 2018; 8(4706):1-5.
34. Abou-shaara H. The origin of honey bee life. *Journal of Entomology and Zoology Studies*. 2015;3(1):239-341.

© 2023 Garba et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/99643>