



## **Antioxidant Activity from *Cymodocea serrulata* Seagrass Crude Extract**

**S. Divyashri<sup>a</sup>, Lakshminarayanan Arivarasu<sup>a\*</sup>, Pitchiah Sivaperumal<sup>b</sup> and Lakshmi Thangavelu<sup>a</sup>**

<sup>a</sup> Department of Pharmacology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai-600077, India.

<sup>b</sup> Department of Pharmacology, Marine Biomedical and Environmental Health Research Laboratory - Blue Lab Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Chennai-600077, India.

### **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i61B35603

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/77780>

**Original Research Article**

**Received 24 October 2021**  
**Accepted 27 December 2021**  
**Published 28 December 2021**

### **ABSTRACT**

**Introduction:** *Cymodocea* can be found in clear water and in the high intertidal areas. The antioxidant study of the extract of *C. serrulata* shows the highest free radical scavenging property on ethanol extract. This may be due to the presence of high phenolic compounds. The study brings out the medicinal value of *C. serrulata* which can be used as a nutraceutical compound in various food and pharmaceutical industries. Antioxidants are defending your body cell from damage caused by free radicals when they accumulated may cause oxidative stress. The synthetic antioxidants are producing more side effects, while natural antioxidants play a major role in scavenging free radicals without side-effects.

**Aim:** To analyse the antioxidant activity from *cymodocea serrulata* sea grass from crude extract.

**Materials and Methodology:** The fresh leaves of *Cymodocea serrulata* were collected from parangipettai coastal area, Tamil Nadu. The seagrass are washed thoroughly with tap water then shade dried on table tissue paper for 2-3 weeks and turn into a fine power. And 10g of dried powdered seagrass samples were mixed with 100ml of methanol/Ethanol.

Using DPPH assay, to extract 0.1ml add equal volume of DPPH 0.1 ml. After 20 min, absorb at 517 nm. Ascorbic acid is used as a standard concentration and DPPH was calculated.

**Results:** The result concludes that *Cymodocea serrulata* showed good antioxidant activity. when concentration increases the percentage of the zone of inhibition is also increased, which shows that it shows a good antioxidant activity.

**Conclusion:** In this study antioxidant activity was checked by using *Cymodocea serrulata*. Using DPPH assay, it has concluded that it has strong antioxidant activity from crude extract. In the future it can be used for medication and further studies can be done by using individual components for various particles.

**Keywords:** Antioxidant activity; *cymodocea serrulate*; sea grass; DPPH assay; marine.

## 1. INTRODUCTION

Sea grass is an endophytic fungus which grows in the southern region. It produces anti metabolite which grows from the marine plant *cymodocea serrulate* [1]. In marine ecosystems, fish depend on the marine plant and vegetation. The *cymodocea* is a ribbon grassy leaf which is a terrestrial and flowering plant which is submerged in the water and it can see more in the subtropical coastal area and it will grow well in muddy and Sandy regions. And it has antibacterial activity. Plant related products are useful in medicinal purposes such as anti inflammatory, anti cancer, antibiotic [2]. Ethanolic leaf shows antioxidant and antibacterial activity. The bioactivity of sea grass is assessed on lung fibroblast cell lines. In this it contains a high level of carotenoids, mainly xanthophylls with antioxidant roles [3]. Endophytic fungi are mostly monocotyledon and dicotyledon and seagrass are monocotyledon and flowering plants [4]. with diphenyl 1- picryl hydrazyl radical it has reduced power on marine extract. Marine algae act as food material and pharmaceutical for treating oxidative disease [5].

Seagrass of the mandapam coast region has high levels of phenol and high reducing power. Antioxidant activity has a high percentage of DPPH radical scavenging activity [6]. And this is used for various oxidative stress related diseases . DPPH activity is better than vitamin -c of the sea grass . and it produces the free radical which has an antioxidant activity [7]. High phenolic and flavonoid content was found in the plant species . and it is found in many herbaceous plants [8]. The antibacterial property of *cymodocea serrulata* was tested against the human pathogen in that ethyl acetate shows maximum activity against the pathogen [9]. It is a potential bio reductant. It grows rapidly and is eco-friendly towards cancer therapy [9,10]. It is more tolerant to burial and light attenuation; it can form a canopy in a higher position above the bottom due to the presence of vertical rhizomes [11].

Chloroform and methanol extract showed effective inhibition against alpha amylase. It shows glucosidase inhibition which is mainly responsible for antidiabetic action [12].

Antioxidant activity of water extract plants was evaluated in that it shows strong antioxidant activity. The plant extracts the highest superoxide radical scavenging activity and acts as a natural antioxidant [12,13]. *Cymodocea spp* used as tranquilizer babies during pregnancy and were used for cough and malaria and leprosy [14]. It shows the reduced MIC and shows no inhibition. In methanol and ethyl acetate it shows maximum pathogen. Natural products produce important sources for antibacterial agents. Marine species comprise total global biodiversity and antimicrobial seagrass encourage travelling organisms. Isolation of sea grass produces biological molecules which prove to produce new drug discovery [15].

There was no study conducted about the antioxidant activity *cymodocea serrulata* in the previous studies. Our team has extensive knowledge and research experience that has translated into high quality publications [16–20],[21-33]. In future, further studies can be done with other activities of *cymodocea serrulata* from different regions and with different activities. And the aim of this study is to determine the antioxidant activity of *cymodocea serrulata* sea grass from crude extract.

## 2. MATERIALS AND METHODOLOGY

### 2.1 Collection of Plant Material and Preparation

The fresh leaves of *Cymodocea serrulata* were collected from Parangipettai coastal area, Tamil Nadu. The seagrass is washed thoroughly with tap water then shade dried on table tissue paper for 2-3 weeks and turned into a fine powder.

## 2.2 Preparation of Extraction

10g of dried powdered seagrass samples were mixed with 100ml of methanol/Ethanol (V/V) and allowed to place for 24 hours at ambient temperature. Then the mixture was passing through whatman filter paper (No.4) then the filtrate was centrifuged at 3000 rpm for 10min and further filtered by 0.45µm syringe micro filter. At last, the solvents are evaporated via vacuum rotary evaporator until samples are obtained in powder form. Then the sample was stored in a shadowy aluminum container at 4°C for further analysis.

## 2.3 Total Antioxidant Activity

Total antioxidant activity of the crude seagrass extract was determined by following method: 0.3 ml of sample was prepared in different concentrations (0.5– 3 mg/ml) with 3 ml of reagent solution (0.6 M sulfuric acid, 28mM sodium phosphate and 4mM ammonium molybdate). Reaction mixture was incubated at 95° c for 90 minutes in a water bath. Absorbance of all sample mixtures was measured at 695 nm. Total antioxidant activity has been expressed as the number of equivalents of ascorbic acid.

## 2.4 DPPH Assay

The antioxidant potential of seagrass crude extract was determined on the basis of their scavenging activity of the stable 1,1- diphenyl-2-picryl hydrazyl (DPPH) free radical. Different concentrations (0.5-3mg/ml) of samples were mixed with 2.9ml diphenylpicrylhydrazyl (DPPH) solution (120µM) in methanol and incubated in darkness at 37°C for 30 minutes. The

absorbance was recorded at 517 nm. Inhibition of free radical by DPPH in percentage (I %) was calculated with the following equation:

$$\text{Percentage of Inhibition (I \%)} = \frac{(A_{\text{blank}} - A_{\text{sample}})}{A_{\text{blank}}} \times 100$$

Where,  $A_{\text{blank}}$  is the absorbance of the control reaction and  $A_{\text{sample}}$  is the absorbance of the test compound. The values of inhibition were calculated for the various concentrations of the sample. Ascorbic acid was used as positive control (Kamala et al., 2015) and all the tests were carried out in triplicate.

## 2.5 Total Reducing Power

Reducing capacity of crude extract obtained from the seagrass extract were determined by following method: Briefly, 1 ml of Benzene: chloroform (2:1) containing different concentrations of extract (0.5-3mg/ml) were mixed with 2.5 ml of benzene: chloroform and 2.5 ml potassium ferricyanide (1%) reaction mixture was incubated at 50° C for 20 min. After incubation, 2.5 ml 10% trichloroacetic acid was added and centrifuged at 10,000 rpm for 10 min 2.5ml. The upper layer was mixed with 2.5ml distilled water and 0.5ml  $\text{FeCl}_3$  (0.1%) and the absorbance was measured at 700 nm.

## 3. RESULTS

In this orange denotes standard solution and blue denotes the percentage of inhibition. When the concentration increases the percentage of inhibition also increases which shows a good antioxidant activity.





Fig.1. The fresh leaves of *Cymodocea serrulata* which are collected from parangipettai coastal area, Tamilnadu



**Fig. 2.** Graph representing the antioxidant activity of *cymodocea serrulata* sea grass. In this x-axis represents the concentration level and y-axis represents the % DPPH inhibition, data implies as mean±SEM

**Table 1.** Total antioxidant activity of *cymodocea serrulata* (sea grass)

TAA	AAE
25 µg/ml	36.85±1.22
50 µg/ml	54.69±1.3
75 µg/ml	72.84±0.9
100 µg/ml	98.61±1.22
125 µg/ml	114.75±1.31
150 µg/ml	137.65±1.3

**Table 2.** Total reducing property of *cymodocea serrulata* sea grass

TRP	AAE
25 µg/ml	12.62±1.21
50 µg/ml	31.49±1.3
75 µg/ml	42.68±0.9
100 µg/ml	57.29±1.23
125 µg/ml	68.38±1.31
150 µg/ml	81.57±1.3

**Table 3.** shows the antioxidant activity of *cymodocea serrulata* by DPPH assay

DPPH	% of inhib	Std
25 µg/ml	19.27	37.3
50 g/ml	41.52	62.7
75 g/ml	54.38	78.52
100 g/ml	67.24	83.59
125 g/ml	78.65	92.4
150 g/ml	94.38	98.6

#### 4. DISCUSSION

In the current study the result shows that *cymodocea serrulata* seagrass show good antioxidant activity from the crude extract. In fig - 2 the graph represents the antioxidant activity of *cymodocea serrulata* sea grass from the crude. In this x-axis represents the concentration level

and y-axis represents the % DPPH inhibition. In this orange denotes standard solution and blue denotes the % of inhibition. When the concentration increases the percentage of inhibition also increases which shows a good antioxidant activity. The antioxidant activity of the plant shows a percentage of 94% which is not as good as the standard but can be used as a

potent antioxidant. Similar studies on the same species show a similar range of antioxidant activity. Previous studies on nutmeg shows that it was an antioxidant of 89.37 percent which is in comparison with the standard of 98.61% doing that these steps can be used as a potent antioxidant [34]. The active extract of *C.serrulata* showed maximum inhibition against *E.coli* and the compound phenyl thioketone was analysed for its antibacterial activity and was proved good and can be used for therapeutic purposes [35,36].

The bioactive compounds from endophytic bacteria showed maximum sensitivity with minimum concentration than the bioactive compounds from epiphytic bacteria and other biological origin. Hence, steps have been undertaken to find out the reason for the maximum activity of endophytic bacteria from seagrasses [37]. In previous studies it is reported that the application of the extract of seagrass proves toxic less and does not produce any harmful effect and it has better free radical scavenging activity which justifies the results of our present study [38-39]. In previous studies it is also reported that the extract has significant Minimum inhibitory concentration and Minimum bacterial concentration against all the bacteria; pathogens and it is more sensitive against *Pseudomonas aeruginosa* [40]. Our team has extensive knowledge and research experience that has translated into high quality publications [41-54].

There are some potential limitations, that is the study is taken into consideration with only one marine plant and it should be considered to be done on a large scale. And there is a high possibility of occurrence of error. The study was carried out in-vitro and it cannot be assumed that the result of antioxidant activity could be translated into clinical effectiveness which proves to be a limitation. In future clinical trials and animal experiments can be done to check the toxicity of the extract. In the future it can be formulated as an alternative drug and commercial products can be prepared which will possess a great and potential value in the herbal markets.

## 5. CONCLUSION

Using DPPH assay, *cymodocea serrulata* sea grass has strong antioxidant activity from crude extract. In the future it can be formulated for medication. It consists of methanolic compounds,

phenolic compounds, flavonoids and isolated components [55-64]. Each of these properties plays a key role in the advancement of human health. From the present study it is evident that Sea grass possesses a good antioxidant activity. The significant health benefits of sea grass have been explored. Further investigations are necessary to provide additional clinical evidence against the free radical scavenging activity.

## DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

It is not applicable.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Supaphon P, Phongpaichit S, Rukachaisirikul V, Sakayaroj J. Antimicrobial potential of endophytic fungi derived from three seagrass species: *Cymodocea serrulata*, *Halophila ovalis* and *Thalassia hemprichii*. PLoS One. 2013 Aug 16;8(8):e72520.
2. Thangaradjou T, Bhatt JR. Status of seagrass ecosystems in India [Internet]. Ocean & Coastal Management. 2018; 159:7–15. Available: <http://dx.doi.org/10.1016/j.ocecoaman.2017.11.025>
3. Sansone C, Galasso C, Lo Martire M, Fernández TV, Musco L, Dell'Anno A, et al. In Vitro Evaluation of Antioxidant Potential of the Invasive Seagrass *Halophila stipulacea* [Internet]. Marine Drugs. 2021;19:37.

- Available:<http://dx.doi.org/10.3390/md19010037>
4. Willette DA, Ambrose RF. Effects of the invasive seagrass *Halophila stipulacea* on the native seagrass, *Syringodium filiforme*, and associated fish and epibiota communities in the Eastern Caribbean [Internet]. *Aquatic Botany*. 2012;103:74–82. Available:<http://dx.doi.org/10.1016/j.aquabot.2012.06.007>
  5. Aljahdali MO, Alhassan AB. Heavy Metal Accumulation and Anti-Oxidative Feedback as a Biomarker in Seagrass *Cymodocea serrulata* [Internet]. *Sustainability*. 2020;12:2841. Available: <http://dx.doi.org/10.3390/su12072841>
  6. Rengasamy RRK, Rajasekaran A, Micheline G-D, Perumal A. Antioxidant activity of seagrasses of the Mandapam coast, India. *Pharm Biol*. 2012 Feb;50(2):182–7.
  7. Hartog CD, Den Hartog C. Seagrasses of coromandel coast, India [Internet]. *Aquatic Botany*. 1994;48:91. Available: [http://dx.doi.org/10.1016/0304-3770\(94\)90075-2](http://dx.doi.org/10.1016/0304-3770(94)90075-2)
  8. Athiperumalsami T, Devi Rajeswari V, Hastha Poorna S, Kumar V, Louis Jesudass L. Antioxidant activity of seagrasses and seaweeds [Internet]. *Botanica Marina*. 2010;53. Available:<http://dx.doi.org/10.1515/bot.2010.032>
  9. Kumar CS, Sarada DVL, Gideon TP, Rengasamy R. Antibacterial activity of three South Indian seagrasses, *Cymodocea serrulata*, *Halophila ovalis* and *Zostera capensis* [Internet]. *World Journal of Microbiology and Biotechnology*. 2008;24:1989–92. Available:<http://dx.doi.org/10.1007/s11274-008-9695-5>
  10. Ravikumar S. Antibacterial activity of *Cymodocea serrulata* root extract against chosen poultry pathogens [Internet]. *Indian Journal of Science and Technology*. 2011;4:98–100. Available:<http://dx.doi.org/10.17485/ijst/2011/v4i2.16>
  11. Buapet P, Makkliang F, Thammakhet-Buranachai C. Photosynthetic activity and photoprotection in green and red leaves of the seagrasses, *Halophila ovalis* and *Cymodocea rotundata*: implications for the photoprotective role of anthocyanin [Internet]. *Marine Biology*. 2017;164. Available:<http://dx.doi.org/10.1007/s00227-017-3215-9>
  12. Unnikrishnan PS, Suthindhiran K, Jayasri MA. Alpha-amylase Inhibition and Antioxidant Activity of Marine Green Algae and its Possible Role in Diabetes Management. *Pharmacogn Mag*. 2015 Oct;11(Suppl 4):S511–5.
  13. Fernando IPS, Kim M, Son K-T, Jeong Y, Jeon Y-J. Antioxidant Activity of Marine Algal Polyphenolic Compounds: A Mechanistic Approach. *J Med Food*. 2016 Jul;19(7):615–28.
  14. Lee J-H, Kim G-H. Evaluation of Antioxidant Activity of Marine Algae-Extracts From Korea [Internet]. *Journal of Aquatic Food Product Technology*. 2015;24:227–40. Available:<http://dx.doi.org/10.1080/10498850.2013.770809>
  15. Lee Y-S, Lee S-H, Kim B-K, Shin K-H. Screening for Aldose Reductase Inhibitory Activity of Extracts of the Marine Plants from Korea [Internet]. *ALGAE*. 2004;19:349–52. Available:<http://dx.doi.org/10.4490/algae.2004.19.4.349>
  16. Rajeshkumar S, Kumar SV, Ramaiah A, Agarwal H, Lakshmi T, Roopan SM. Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells. *Enzyme Microb Technol*. 2018 Oct;117:91–5.
  17. Nandhini NT, Rajeshkumar S, Mythili S. The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation. *Biocatal Agric Biotechnol*. 2019 May 1;19:101138.
  18. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of *Enterococcus* sp.–mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. *Environ Sci Pollut Res*. 2020 Mar 1;27(8):8166–75.
  19. Gomathi M, Prakasam A, Rajkumar PV, Rajeshkumar S, Chandrasekaran R, Anbarasan PM. Green synthesis of silver nanoparticles using *Gymnema sylvestre* leaf extract and evaluation of its antibacterial activity [Internet]. *South African Journal of Chemical Engineering*. 2020;32:1–4. Available:<http://dx.doi.org/10.1016/j.sajce.2019.11.005>

20. Rajasekaran S, Damodharan D, Gopal K, Rajesh Kumar B, De Poures MV. Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends. *Fuel*. 2020 Oct 1;277:118166.
21. Santhoshkumar J, Sowmya B, Venkat Kumar S, Rajeshkumar S. Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*. *S Afr J Chem Eng*. 2019 Jul;29:17–23.
22. Raj RK, DE, SR.  $\beta$ -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line. *J Biomed Mater Res A*. 2020 Sep;108(9):1899–908.
23. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb Pathog*. 2018 Apr;117:68–72.
24. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells. *Hum Exp Toxicol*. 2019 Jun 1;38(6):694–702.
25. Ezhilarasan D, Sokal E, Najimi M. Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets. *Hepatobiliary Pancreat Dis Int*. 2018 Jun;17(3):192–7.
26. Ezhilarasan D. Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective. *Arab J Gastroenterol*. 2018 Jun;19(2):56–64.
27. Gomathi AC, Xavier Rajarathinam SR, Mohammed Sadiq A, Rajeshkumar S. Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of *Tamarindus indica* on MCF-7 human breast cancer cell line. *J Drug Deliv Sci Technol*. 2020 Feb 1;55:101376.
28. Dua K, Wadhwa R, Singhvi G, Rapalli V, Shukla SD, Shastri MD, et al. The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress. *Drug Dev Res*. 2019 Sep;80(6):714–30.
29. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol*. 2018 Oct;89(10):1241–8.
30. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol*. 2021 Feb;122:105030.
31. Joseph B, Prasanth CS. Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry? *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2021 Jul;132(1):118–9.
32. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. *Syzygium cumini* extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med*. 2019 Feb;48(2):115–21.
33. Duraisamy R, Krishnan CS, Ramasubramanian H, Sampathkumar J, Mariappan S, Navarasampatti Sivaprakasam A. Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments. *Implant Dent*. 2019 Jun;28(3):289–95.
34. Charles DJ. *Antioxidant Properties of Spices, Herbs and Other Sources*. Springer Science & Business Media; 2012:612.
35. Sivaperumal P, Kamala K, Rajaram R. Biosorption of Long Half-life Radionuclide of Strontium Ion (Sr) by Marine Actinobacterium *Nocardiostrictum* sp. 13H [Internet]. *Geomicrobiology Journal*. 2018;35:300–10. Available:<http://dx.doi.org/10.1080/01490451.2017.1350891>
36. Kamala K, Sivaperumal P, Gobalakrishnan R, Swarnakumar NS, Rajaram R. Isolation and characterization of biologically active alkaloids from marine actinobacteria *Nocardiostrictum* sp. NCS1 [Internet]. *Biocatalysis and Agricultural Biotechnology*. 2015;4:63–9. Available:<http://dx.doi.org/10.1016/j.bcab.2014.10.005>
37. Sivaperumal P, Kamala K, Rajaram R. Bioremediation of Industrial Waste Through Enzyme Producing Marine Microorganisms [Internet]. *Marine Enzymes Biotechnology: Production and Industrial Applications, Part III - Application of Marine Enzymes*. 2017: 165–79. Available:<http://dx.doi.org/10.1016/bs.afnr.2016.10.006>

38. Brahmachari G. Biotechnology of Microbial Enzymes: Production, Biocatalysis and Industrial Applications. Academic Press. 2016:632.
39. Ananthan G, Sivaperuma P, Mohamed Hussain S. Antibacterial Potential of Marine Ascidian *Phallusia arabica* Against Isolated Urinary Tract Infections Bacterial Pathogens [Internet]. Asian Journal of Animal Sciences. 2011;5:208–12. Available:<http://dx.doi.org/10.3923/ajas.2011.208.212>
40. Darshit R, Pandya D. Screening and characteristic study of antimicrobial actinomycetes from near-by soil of medicinal plants [Internet]. International Journal of Pharmacy and Pharmaceutical Sciences. 2018;10:66. Available:<http://dx.doi.org/10.22159/ijpps.2018v10i11.29068>
41. Pushpaanjali G, Geetha RV, Lakshmi T. Knowledge and Awareness about Antibiotic Usage and Emerging Drug Resistance Bacteria among Dental Students. Journal of Pharmaceutical Research International. 2020: 34–42.
42. Aathira CM, Geetha RV, Lakshmi T. Knowledge and Awareness about the Mode of Transmission of Vector Borne Diseases among General Public. Journal of Pharmaceutical Research International. 2020 Aug 24;87–96.
43. Baskar K, Lakshmi T. Knowledge, Attitude and Practices Regarding HPV Vaccination among Undergraduate and Postgraduate Dental Students in Chennai. Journal of Pharmaceutical Research International. 2020 Aug 25;95–100.
44. Manya Suresh LT. Wound Healing Properties of Aloe Barbadensis Miller-In Vitro Assay. Journal of Complementary Medicine Research. 2020;11(5):30–4.
45. First Report on Marine Actinobacterial Diversity around Madras Atomic Power Station (MAPS), India [Internet]. [Cited 2021 Aug 31]. Available:<http://alinteridergisi.com/article/first-report-on-marine-actinobacterial-diversity-around-madras-atomic-power-station-maps-india/>
46. Physicochemical Profile of Acacia Catechu Bark Extract – An in Vitro Stud - International Journal of Pharmaceutical and Phytopharmacological Research [Internet]. [Cited 2021 Aug 31]. Available:<https://eijppr.com/article/physicochemical-profile-of-acacia-catechu-bark-extract-an-in-vitro-stud>
47. Lakshmi T. Antifungal Activity of *Ficus racemosa* Ethanolic Extract against Dermatophytes-An in vitro Study. Journal of Research in Medical and Dental Science. 2021;9(2):191–3.
48. Awareness of Drug Abuse among Teenagers - International Journal of Pharmaceutical and Phytopharmacological Research [Internet]. [Cited 2021 Aug 31]. Available:<https://eijppr.com/article/awareness-of-drug-abuse-among-teenagers>
49. Mangal CSK, Anitha R, Lakshmi T. Inhibition of Nitric oxide Production and Nitric oxide Synthase Gene Expression in LPS Activated RAW 264 .7 Macrophages by Thyme oleoresin from *Thymus vulgaris*. J Young Pharm. 2018;10(4):481.
50. COX2 Inhibitory Activity of *Abutilon Indicum* - Pharmaceutical Research and Allied Sciences [Internet]. [Cited 2021 Aug 31]. Available: <https://ijpras.com/article/cox2-inhibitory-activity-of-abutilon-indicum>
51. Jibu RM, Geetha RV, Lakshmi T. Isolation, Detection and Molecular Characterization of *Staphylococcus aureus* from Postoperative Infections. Journal of Pharmaceutical Research International. 2020 Aug 24;63–7.
52. Sindhu PK, Thangavelu L, Geetha RV, Rajeshkumar S, Raghunandhakumar S, Roy A. Anorectic drugs: an experimental and clinical perspective -A Review. Journal of Complementary Medicine Research. 2020;11(5):106–12.
53. Nivethitha R, Thangavelu L, Geetha RV, Anitha R, RajeshKumar S, Raghunandhakumar S. In Vitro Anticancer Effect of *Sesamum Indicum* Extract -. Journal of Complementary Medicine Research. 2020;11(5):99–105.
54. Mariona P, Roy A, Lakshmi T. Survey on lifestyle and food habits of patients with PCOS and obesity. Journal of Complementary Medicine Research. 2020;11(5):93–8.
55. Rajendran R, Kunjusankaran RN, Sandhya R, Anilkumar A, Santhosh R, Patil SR. Comparative Evaluation of Remineralizing Potential of a Paste Containing Bioactive Glass and a Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate: An in Vitro Study.



- Pesqui Bras Odontopediatria Clin Integr. 2019 Mar 12;19(0):4668.
56. Ashok BS, Ajith TA, Sivanesan S. Hypoxia-inducible factors as neuroprotective agent in Alzheimer's disease. Clin Exp Pharmacol Physiol [Internet]. 2017 Mar [cited 2021 Sep 15];44(3). Available: <https://pubmed.ncbi.nlm.nih.gov/28004401/>
57. Malli SN, Selvarasu K, Jk V, Nandakumar M, Selvam D. Concentrated Growth Factors as an Ingenious Biomaterial in Regeneration of Bony Defects after Periapical Surgery: A Report of Two Cases. Case Rep Dent [Internet]. 2019 Jan 22 [Cited 2021 Sep 15];2019. Available: <https://pubmed.ncbi.nlm.nih.gov/30805222/>
58. Mohan M, Jagannathan N. Oral field cancerization: an update on current concepts. Oncol Rev [Internet]. 2014 Jun 30 [Cited 2021 Sep 15];8(1). Available: <https://pubmed.ncbi.nlm.nih.gov/25992232/>
59. Menon S, Ks SD, R S, S R, Vk S. Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism. Colloids Surf B Biointerfaces [Internet]. 2018 Oct 1 [cited 2021 Sep 15];170. Available: <https://pubmed.ncbi.nlm.nih.gov/29936381/>
60. Samuel SR, Acharya S, Rao JC. School Interventions-based Prevention of Early-Childhood Caries among 3-5-year-old children from very low socioeconomic status: Two-year randomized trial. J Public Health Dent [Internet]. 2020 Jan [cited 2021 Sep 15];80(1). Available: <https://pubmed.ncbi.nlm.nih.gov/31710096/>
61. Praveen K, Narayanan V, Muthusekhar MR, Baig MF. Hypotensive anaesthesia and blood loss in orthognathic surgery: a clinical study. Br J Oral Maxillofac Surg [Internet]. 2001 Apr [cited 2021 Sep 15];39(2). Available: <https://pubmed.ncbi.nlm.nih.gov/11286449/>
62. Neelakantan P, Subbarao C, Subbarao CV, De-Deus G, Zehnder M. The impact of root dentine conditioning on sealing ability and push-out bond strength of an epoxy resin root canal sealer. Int Endod J [Internet]. 2011 Jun [cited 2021 Sep 15];44(6). Available: <https://pubmed.ncbi.nlm.nih.gov/21255047/>
63. Oligonucleotide therapy: An emerging focus area for drug delivery in chronic inflammatory respiratory diseases. Chem Biol Interact. 2019 Aug 1;308: 206–15.
64. Kumar MS, Vamsi G, Sripriya R, Sehgal PK. Expression of matrix metalloproteinases (MMP-8 and -9) in chronic periodontitis patients with and without diabetes mellitus. J Periodontol. 2006 Nov;77(11):1803–8.

© 2021 Divyashri et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/77780>