



Analysis of HIV/Malaria Coinfections among HIV-1 Infected Individuals in Two Tertiary Hospitals in Old Cross River State, Nigeria

**Ugochi Immaculate Ejike¹, Tochi Ifeoma Cookey¹,
Hope Chioma Innocent-Adiele² and Iheanyi Omezuruike Okonko^{1*}**

¹*Virus Research Unit, Department of Microbiology, University of Port Harcourt, Port Harcourt, Nigeria.*
²*Department of Applied Microbiology, Ebonyi State University, Abakaliki, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author IOO conceived and designed the study, partially financed the whole study, assisted in the collection of samples, the laboratory analysis and contributed principally in writing up the manuscript as well as in the data and statistical analysis of this study. Author UIE procured the materials used, ran the laboratory analysis and drafting of the manuscript which she used as part of her M.Sc. thesis in the Department of Microbiology, University of Port Harcourt, Port Harcourt, Nigeria. Author HCIA wrote the proposal for ethical approval, procured the materials used for sample collection, addressed the subjects, collected samples and participated in the study design and reviewing of the manuscript. Author TIC participated in the laboratory analysis and performed the statistical analysis. All authors contributed equally in writing up this manuscript, editing, proof-reading and approval of the final manuscript for this publication.

Article Information

DOI: 10.9734/IJTDH/2020/v41i1030327

Editor(s):

(1) Dr. Shankar Srinivasan, Rutgers School of Health Related Professions, United States of America.

Reviewers:

(1) Ikenna Daniel Molobe, Unified Initiative for a Drug-Free Nigeria, Lagos, Nigeria.

(2) Olusimbo Ige, General Board of Global Ministries, United States of America.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/57414>

Original Research Article

Received 19 May 2020
Accepted 24 July 2020
Published 05 August 2020

ABSTRACT

Aims: Human Immunodeficiency Virus (HIV) and Malaria are the two main global public health threats that dent development in low and middle-income countries. This study evaluated the immunological marker and HIV/Malaria co-infection among individuals infected with HIV-1 in old Cross River State, Nigeria.

Study Design: Cross-sectional study.

Place and Duration of Study: University of Calabar Teaching Hospital (UCTH) and University of Uyo Teaching Hospital (UUTH) between March 2018 and August 2019.

*Corresponding author: Email: iheanyi.okonko@uniport.edu.ng;

Methods: A total of 417 individuals infected with HIV-1 partook in this study. The age of these individuals ranged from 4-72 years (average age = 39.1 years). Plasma samples were analyzed for HIV and Malaria using Enzyme-Linked immunosorbent Assay. The CD4 count was enumerated using the Partec CyFlow[®] Counter. Plasma viral loads (PVL) were determined using the Abbott Real-Time HIV-1 assay.

Results: Results showed that 230(55.1%) of the participants were in the 31-45 years age range. The majority (67.4%) of the HIV-1 infected individuals were females. An overall prevalence of HIV/Malaria coinfection in Old Cross River State, Nigeria was 14.3%, of which Uyo was 6.3% and Calabar was 3.0%. A higher prevalence of HIV/Malaria coinfection was observed among age groups <25 years (17.5%), males (5.1%), singles or divorced/widow/widower (7.7%), those with primary education (7.5%), and students (10.0%). Higher HIV/Malaria coinfection was also observed among those with CD4 cell count <200 cells/ μ l and 350-499 cells/ μ l (5.7%) and PVL >5000 copies/mL (7.9%) compared to others with 2.0% prevalence. Of all variables evaluated only marital status ($p= 0.033$), educational background ($p= 0.000$) and occupations ($p =0.000$) were significantly associated.

Conclusions: This study further confirmed the presence of HIV/Malaria coinfection in old Cross River State, Nigeria. This study has added to the voices on the ground to give a better view on the frequency and the pattern of distribution of HIV/Malaria coinfection since limited studies have been done on this in old Cross River State, Nigeria. This, therefore, highlights the need for a well-structured approach to the management of HIV/Malaria coinfection in Nigeria.

Keywords: Antibodies; HIV; coinfections; malaria; Nigeria.

1. INTRODUCTION

Human Immunodeficiency Virus (HIV) and malaria are two main worldwide health threats that affect development in low and middle-income countries [1-2]. In sub-Saharan Africa, HIV and AIDS coinfection with Malaria is very much prevalent [3]. These infections have led to the loss of about 5 million lives yearly, bearing significant humanitarian, economic, and social impact, which is yet to be measured [1-2]. In addition to occurring in the same zone geographically, these infections are also said to be linked with poverty, having a major public health implication [4].

It is a well-known opportunistic infection and also an important factor affecting the increased disease progression to AIDS in HIV-infected persons due to a significantly increased destruction of active CD4 helper T-cells and a quick breakdown of immune functions. *P. falciparum* causes the severe form of malaria while a milder disease which is less severe is caused by *P. vivax*, *P. ovale*, and *P. malariae* [5]. Over 50% of the people on earth are vulnerable to the malaria infection and they have estimated over 300 million malaria cases annually in the tropics. In 2016, Nigeria contributed to over 27% of the total malaria cases and 24% of mortality from malaria globally. Some studies have revealed a high occurrence of malaria among people living with HIV [6]. A population-based study showed that those infected with HIV were

more likely to get Malaria than those that were not infected with the virus [7].

The high occurrence of both malaria and HIV in the sub-Saharan region indicates that coinfection is relatively common, making it an important health concern due to the synergistic nature of the HIV-malaria partnership. HIV infection exerts influence on malaria by diminishing the response of the immune system to Plasmodium invasion and it's been maintained by several researchers that malaria increases the replication of the HIV both *in vivo* and *in vitro* [6].

Sub-Saharan Africa is hit by most deadly diseases in the world. HIV and malaria are diseases that mostly affect the poor, contributing greatly to the high poverty level in sub-Saharan Africa nations. The young population whose strength contribute to the growth of the local economy are mostly hit by these diseases [8]. The risk and gravity of Malaria disease are amplified by HIV. Infections with HIV and malaria are amongst the two most vital global health problems of developing countries including Nigeria, which was said to cause more than 4 million deaths a year [9].

Furthermore, the rate at which malaria is spread is boosted when there is HIV infection. This process activates CD4 cells and increases the production of cytokines which makes the microenvironment suitable for the distribution of HIV among the CD4 cells, thereby increasing

HIV-1 replication. Researches done previously has tried to create a relationship between HIV and malaria, but with difficulties [10]. However, recent researches have proposed that frequent malaria infection culminates in a much faster decline in CD4+ T cells over time. Malaria co-infection with HIV produces regular occurrence of symptomatic malaria [11] and more occurrence of serious or complicated malaria including death in both children and adults [12].

Since 2009, the Centre for Disease Prevention and Control has included Malaria in the list of AIDS-related opportunistic infections. Although not regarded as the major cause of death, malaria ranks third as the most significant cause of ill-health in people living with HIV/ AIDS [13]. The synergistic nature of the HIV/malaria partnership makes coinfection a major health concern. Malaria is said to upsurge HIV replication in vitro and in vivo [11]. Additionally, evidence shows that malaria coinfection with HIV triggers the progression of malaria disease, enhances the risk of severe malaria in adults [14] and raises the risk of congenital infection [15]. Furthermore, this dual infection increases the distribution of both diseases, particularly in sub-Saharan Africa [12,16-18].

A previous study [19] investigating the frequency of HIV and Malaria seropositivity amongst undergraduates at a tertiary healthcare Centre in Port Harcourt, Nigeria showed no incidence of coinfection. Consequently, the present study aimed at determining the prevalence of HIV/Malaria coinfections in HIV-1 infected individuals in old Cross River State, Nigeria.

2. MATERIALS AND METHODS

2.1 Study Areas

The study was conducted in the old Cross River States (Akwa Ibom State and Cross Rivers State). Akwa Ibom is one of Nigeria's 36 states, with a population of over five million people created in 1987 from the former Cross River State and is currently the highest oil- and gas-producing state in the country [20]. The state's capital is Uyo, with over 500,000 inhabitants. It is located in the coastal southern part of the country, lying between latitudes 4°32'N and 5°33'N, and longitudes 7°25'E and 8°25'E. The state is located in the South-South geopolitical zone and is bordered on the east by Cross River State, on the west by Rivers State and Abia State, and the south by the Atlantic Ocean and the southernmost tip of Cross River State

[20]. In addition to English, the main spoken languages are Ibibio, Annang, Eketand Oron [21]. Cross River State is a coastal state in southeastern Nigeria [22], bordering Cameroon to the east. It is located in Nigeria's Delta Region and occupies 20,156 square kilometres. It is located at an elevation of 152 meters above sea level and its population amounts to 3,104,446. Its coordinates are 5°45'0" N and 8°30'0" E in DMS (Degrees Minutes Seconds) or 5.75 and 8.5 (in decimal degrees). It shares boundaries with Benue State to the north, Ebonyi and Abia States to the west, to the east by the Cameroon Republic and to the south by Akwa-Ibom and the Atlantic Ocean [23]. Its capital is at Calabar, and it is named after the Cross River [24], which passes through the state. The state is ethnically diverse, including Efiks, Bekwara, and Ejagham inhabitants. The Latitude of Cross River State is 6.167 and the Longitude is 8.6601. Cross River State is connected with a major highway to its sister state Akwa Ibom [20].

2.2 Study Design

A cross-sectional study in old Cross Rivers State, Nigeria was carried out.

2.3 Sample Size Determination

The sample size for this study was determined using the formula: $N = Z^2PQ/d^2$ [25]. Where N is the desired sample size, P is the expected prevalence in the target population, Q = 1-P, Z = 1.96 standard error, d is the level of statistical significance (0.05).

For Akwa-Ibom State, A p value of 6.4% (reported by UNAIDS [26]) was used.

Z= Normal standard distribution that corresponds to 95.0% confidence interval as 1.96.

p= Prevalence of HIV (6.4 % for Akwa-Ibom State as at UNAIDS [26]) = 0.064

q=1-p (1-0.064) = 0.936

d=degree of accuracy/precision expected as 0.05

N= 92 HIV HIV-positive persons (estimated for Akwa-Ibom State).

Hence, the estimated sample size was 92 HIV-positive individuals with an additional 10% sampled to take care of study participants that may be lost to follow-up [25,27], providing a total sample size of 100 for Akwa-Ibom State, Nigeria.

For Cross River State, A p value of 4.4% (reported by UNAIDS, 2016) was used.

Z= Normal standard distribution that corresponds to 95.0% confidence interval as 1.96.

p= Prevalence of HIV (4.4% Cross Rivers State as at UNAIDS [26] sentinel survey) = 0.044

q=1-p (1-0.044) = 0.956

d=degree of accuracy/precision expected as 0.05

N= 65 HIV-positive persons (estimated for Cross Rivers State).

Hence, the estimated sample size was 65 HIV-positive individuals with an additional 10.0 % sampled to take care of study participants that may be lost to follow-up [25,27], providing a total sample size of 75 for Cross River State, Nigeria. However, in this study, in all 417 samples were obtained from all confirmed HIV-1 infected participants. The 417 samples of these cohorts were used for malaria analysis.

2.4 Study Population

Four hundred and seventeen (417) HIV-positive patients were selected and enrolled for the study (Table 1). While the entire HIV- positive individuals in old Cross River State were the target population to which the findings of the study was extrapolated. Their ages ranged from 4-72 years (average = 39.1 years). Two hundred and thirty (55.1%) of them were in the 31-45 years age range. The majority (67.4%) of the HIV-1 infected individuals was females and 32.6% were males. The male: female ratio in the present study of 1: 2 (Table 1). Over 60.0% of the HIV-1 infected individuals were married (62.8%, n=262), 142 (34.1%) were singles and 13(3.1%) were divorced/widows/widowers (Table 1). Majority of the study participants had tertiary education (54.4%, n=227), 147 (35.3%) had secondary education, 40(9.6%) had primary education and 3 (0.7%) had no formal education (Table 1). In terms of occupation, the majority were traders (23.3%), followed by civil servants (15.6%), students (14.4%), business (11.8%) and teaching was 9.6%. About 6.5% were unemployed, 6.0% were artisans, driving and retirees were 3.4% respectively while farmers (2.0%) were the least (Table 1).

2.4.1 Sample collection

Blood samples were collected from HIV-infected individuals in old Cross River State, Nigeria used

in this study. Three millilitres (3 ml) of blood were collected from 417. HIV-infected individuals by venipuncture, between June 2018 and August 2019. The blood was allowed to clot and centrifuged at 3000 rpm for 5 minutes. The sera were carefully aspirated into plain bottles and stored at -20°C until analyzed.

2.5 Serological Analysis of Malaria

Plasma samples were analyzed for the presence of Malaria *Plasmodium falciparum* using the ELISA kit manufactured by DIA.PRO Diagnostic Bioprobes Srl (Milano) – Italy, according to manufacturer's specifications.

2.6 CD4 T Cell Count Enumeration

EDTA-treated blood samples were used for CD4 cell count using Partec CyFlow® Counter (Partec GmbH, Munster, Germany). The CD4 cell count was done as stipulated by the Partec CyFlow® Counter manufacturer.

2.7 HIV-1 Viral Load Testing (Abbott Real-Time Assay)

Plasma viral load (PVL) was analyzed using the Abbott Real-Time HIV assay US Protocol.

2.8 Data Analysis

Descriptive statistics were used to describe the characteristics of the study participants. Chi-square test and a multivariate test (logistic regression) were used to fine-tune our findings and to determine relationships between the variables and the prevalence of HIV/Malaria coinfections. P-value was considered significant at 0.05. All analyses were conducted using complex samples analysis of the Statistical Package for the Social Sciences (SPSS), IBM version 22.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 General characteristics of HIV-infected individuals

Table 1 shows the 417 HIV-infected individuals who partook in this study; 241 (57.8%) from Calabar, Cross Rivers State, Nigeria and 176 (42.2%) from Uyo, Akwa-Ibom State, Nigeria. Clinical characteristics of HIV-infected individuals revealed that the CD4 (cells/ μ l) count ranged

from 5 – 2139 cells/ μ l (average = 455.6 cells/ μ l) (Table 1). Generally, of the 417 samples analyzed for viral loads (VL), 151 (36.2%) had VL less than 40 copies/mL, 102 (24.5%) fell within the range 40-5000 copies/mL, and 164 (39.3%) were within >5001 copies/mL (Table 1).

Table 1. Socio-demographical and clinical characteristics of HIV-1/malaria coinfected individuals in old cross State, Nigeria

Variables	No. tested (%)	Malaria (%)	P value	
Locations				
Uyo	176(58.0)	11(6.3)	P= 0.232	
Calabar	241 (42.0)	7(3.0)		
Age groups (years)				
\leq 25	40(9.6)	7(17.5)	P=0.899	
26-30	46(11.0)	2(4.3)		
31-35	64(15.3)	1(1.6)		
36-40	94(21.8)	1(1.1)		
41-45	72(17.3)	0(0.0)		
46-50	43(10.3)	1(2.3)		
51-55	28(6.7)	1(3.6)		
56-60	21(5.0)	1(4.8)		
\geq 61	12(2.9)	1(8.3)		
Sex				
Males	136(32.6)	7(5.1)	P=0.910	
Females	281(67.4)	11(3.9)		
Marital status				
Married	262(62.8)	6(2.3)	P=0.033	
Singles	142(34.1)	11(7.7)		
Divorced/Widowed	13(3.1)	1(7.7)		
Educational status				
Non-Formal	3(0.7)	0(0.0)	P=0.000	
Primary	40(9.6)	3(7.5)		
Secondary	147(35.3)	7(4.8)		
Tertiary	227(54.4)	8(3.5)		
Occupation				
Trading	97(23.3)	5(5.2)	P=0.000	
Teaching	40(9.6)	1(2.5)		
Civil Servant	65(15.6)	2(3.1)		
Public Servant	18(4.3)	0(0.0)		
Business	49(11.8)	3(6.1)		
Artisans	25(6.0)	0(0.0)		
Driving	14(3.4)	1(7.1)		
Retired	14(3.4)	0(0.0)		
Farming	8(2.0)	0(0.0)		
Student	60(14.4)	6(10.0)		
Unemployed	27(6.5)	0(0.0)		
CD4 counts (cells/μl)				
< 200	70(16.8)	4(5.7)		P=0.758
200-349	103(24.7)	5(4.9)		
350-499	88(21.1)	5(5.7)		
\geq 500	156(37.4)	4(2.6)		
Viral load (copies/mL)				
< 40	151(36.2)	3(2.0)	P=0.120	
40 – 5000	102(24.5)	2(2.0)		
5001 & above	164(39.3)	13(7.9)		
Total	417(100.0)	18(4.3)		

3.1.2 Overall prevalence of HIV/Malaria coinfections

Results showed an overall prevalence of HIV/Malaria coinfection was 4.3% for Old Cross River State, Nigeria, 6.3% for Uyo and 3.0% for Calabar, Nigeria. However, these differences were not statistically associated ($P=0.232$). Table 1 shows the seroprevalence of HIV coinfections with Malaria amongst HIV-1 infected individuals in old Cross Rivers State, Nigeria with their sociodemographic and clinical variables.

3.1.3 Age-specific HIV/Malaria coinfections

Higher HIV/Malaria coinfection was observed among age groups <25 years (17.5%) than in other age-groups. The age-specific prevalence showed that HIV/Malaria coinfections were highest in ages <25 years (17.5%), followed by ages 61 years and above (8.3%), 56-60 years (4.8%), 26-30 years (4.3%), 51-55 years (3.6%), 46-50 years (2.3%) and 31-35 years (1.6%) while 36-40 years had the least prevalence (1.1%), however, these differences were not statistically associated ($P=0.899$) [Table 1].

3.1.4 Sex-specific HIV/Malaria coinfections

Higher HIV/Malaria coinfection was observed among males (5.1%) than in females (3.9%). The study showed no significant difference ($P=0.910$) between sex and HIV/Malaria coinfections amongst HIV-1 infected individuals (Table 1).

3.1.5 Marital status-specific HIV/Malaria coinfections

Higher HIV/Malaria coinfection was observed among individuals who were singles or divorced/widow/widower (7.7%) than in married individuals (2.3%). A significant difference ($P=0.033$) exist between marital status and HIV/Malaria coinfections amongst HIV-1 infected individuals (Table 1).

3.1.6 Educational status-specific HIV/Malaria coinfections

Higher HIV/Malaria coinfection was observed among individuals who had primary education (7.5%) than other educational status (Secondary 4.8%, Tertiary 3.5%). A significant difference ($P=0.000$) exist between educational status and HIV/Malaria coinfections (Table 1).

3.1.7 Occupation-specific HIV/Malaria coinfections

Higher HIV/Malaria coinfection was observed among students (10.0%) than other occupations. This was followed by drivers (7.1%), business (6.1%), traders (5.2%), civil servants (3.1%) and teachers (2.5%) while other occupations recorded zero prevalence for HIV/Malaria coinfection. A significant difference ($P=0.000$) exist between occupation and HIV/Malaria coinfections (Table 1).

3.1.8 CD4-specific HIV/Malaria coinfections

Higher HIV/Malaria coinfection was observed among HIV-1 patients with CD4 T cell count <200 cells/ μ l and 350-499 cells/ μ l (5.7%), followed by those with CD4 T cell count 200-349 cells/ μ l (4.9%) and the least prevalence occurred in those with CD4 T cell count >500 cells/ μ l (2.6%) as shown in Table 1. These differences were not statistically significant ($P=0.758$).

3.1.9 Viral loads-specific HIV/Malaria coinfections

Higher HIV/Malaria coinfection was observed among HIV-1 patients with viral loads (VL) above 5000 copies/mL (7.9%) compared to others with 2.0% prevalence (Table 1). No significant difference ($P=0.120$) exist between Viral load and HIV/Malaria coinfections (Table 1).

3.2 Discussion

Malaria is still a major public health problem especially within the tropical and sub-tropical regions despite a clear decline in its incidence by 30% globally and by 34% in Africa [28-30]. The need to conduct prevalence studies and surveys such as this on an annual basis is to monitor the progress made so far in realizing Sustainable Development Goal (SDG) of combating malaria and HIV [28]. It is also in a way to measure the impact and effectiveness of malaria control programmes in the area [28]. Evidence-based studies in other states in Nigeria have documented a high prevalence of malaria among pregnant women such as 26% in Port Harcourt [31], 40.6% in Abakaliki [32], 40.7% in Cross River State [28], 52.0% reported in Lagos [33] and 99.0% in Enugu [34]. In the same vein, high prevalences as high as 48.06% to 70.8% were also reported among children under five years of age in other previous studies in Nigeria [28,35-38]. Malaria prevalence was reported to be 48.06% in Jos [35], 58.2% in Anambra [36], 64%

in Abuja [37], 66.3% in Cross River State [28] and 70.8% in Ogun [38]. Previous studies in old Cross River State [39-40] had reported as high as 55.0% and 80.8% prevalence of malaria respectively, in the general population in Cross River State, Nigeria. In the same vein, 12.0% prevalence of malaria was reported in the general population in Akwa Ibom State, Nigeria. Also, in the general population, previous studies [41,42] had reported a total of 3,228 cases of malaria from 1983–2012, with a minimum of 61 and maximum cases of 302. Since these evidenced-based studies in general populations had reported high malaria transmission, it was therefore pertinent to carry out a coinfection study among the HIV population in the settings. It was based on this evidence that this study was aimed at determining the prevalence of HIV/malaria co-infection in old Cross River State, Nigeria.

This study showed that HIV coinfections with Malaria were determined to be 4.3%. The 4.3% coinfection rate reported in this study is dissimilar when compared with the findings of other studies elsewhere in Nigeria. Dada et al. [43] in a study carried out in Akure, Ondo State, Nigeria reported HIV/Malaria coinfection to be 10.3%. This low HIV/Malaria coinfection (4.3%) observed in this study is similar to a study carried out among undergraduate students in Port Harcourt with zero prevalence [19]. A similar value (4.55%) was reported in Akure [44]. Another similar study carried out in Bamenda Cameroon also indicated a low coinfection rate (2.24%) of HIV/malaria [45]. Olusi and Abe [44] carried out a study which was intended to investigate the level of coinfection of HIV and Malaria parasite in Akure, Ondo State, Nigeria, that study revealed that of 4.55% of the pregnant women tested, was infected with HIV and 96.92% were infected with malaria.

Higher HIV/Malaria coinfection was observed among age groups ≤ 25 years (17.5%) than in other age-groups. Similarly, higher HIV/Malaria coinfection was also observed among age groups ≤ 25 years (10.0%) than in other age-groups. This is slightly different from a study where higher HIV/Malaria coinfection was observed in ages of 20-49 years [43].

Gender has been highlighted as an important risk factor in the frequency of both malaria and HIV disease with women being 50% more at risk of contracting malaria than men [46]. Higher HIV/Malaria coinfection was observed among males (5.1%) than in females (3.9%). Though,

the difference was not significant ($P > 0.05$). This finding is similar to another study in Akure, Ondo State, Nigeria which reported HIV/Malaria coinfection to be 10.3% with a higher frequency in females than in males [43]. Also, similar to a study carried out in Bamenda Cameroon of which 6 out of 7 that were coinfecting with Malaria were females [44]. In other studies, in old Cross River State, Nigeria [41-42], there were more cases of malaria in males than in females. This simply implies that males are more infected by malaria than their female counterparts probably due to their immunological characteristics [41-42].

Higher HIV/Malaria coinfection was observed among individuals who were singles or divorced/widow/widower (7.7%) than in married individuals (2.3%). Higher HIV/Malaria coinfection was observed among individuals who had primary education (7.5%) than other educational status (Secondary 4.8%, Tertiary 3.5%). This is contrary to many studies which have shown the frequency of coinfection to be higher among those with no formal education [47]. A significant difference ($P < 0.05$) exist between occupation and HIV/Malaria coinfections.

HIV-infected individuals are at an increased risk of clinical malaria, severe illness, hospitalization, and death [48]. Malaria causes anaemia leading to blood transfusions, a procedure that increases the risk for HIV infection where universal blood screening is yet to be achieved [48]. From this standpoint, our study reported higher HIV/Malaria coinfection was observed among HIV-1 patients with CD4 cell count < 200 cells/ μ l and 350-499 cells/ μ l (5.7%), followed by those with CD4 T cell count 200-349 cells/ μ l (4.9%) and the least prevalence occurred in those with CD4 cell count ≥ 500 cells/ μ l (2.6%). In a related study by Eyong et al. [3], lower CD4 count levels were significantly related to high-density malaria parasitemia among in HIV and AIDS individuals in Bamenda, Cameroon.

Malaria contributes to a temporary increase in viral load among HIV-infected people which may worsen the clinical disease, increase mother-to-child transmission, and augment transmission in adults [48]. Also, based on this premise, 39.3% of the individuals with HIV-1 in this study had a high viral load above 5,000 copies/ml which is indicative of treatment failure, defined by WHO [49] as a persistently detectable viral load exceeding 1000 copies/ml (that is, two consecutive viral load measurements within a

three-month interval, with adherence support between measurements) after at least six months of using ARV drugs. Higher HIV/Malaria coinfection was observed among HIV-1 patients with viral loads (VL) above 5000 copies/mL (7.9%) compared to others with 2.0% prevalence. An important study from Malawi revealed that HIV-1 plasma viral loads were high in malaria-infected individuals than in those not infected with Malaria, and these levels remained higher for up to 10 weeks after treatment [50].

From evidence-based studies in the study area, the trend in malaria prevalence from 1983 to 2012 revealed a 75.5% increase in malaria in old Cross River State, Nigeria. This is owing to the reported cases of malaria reported from 2006-2012 were more significant than those reported in the 1980s [41,42]. The pattern obtained in their study indicates that malaria is still on the increase as it has not been properly tackled in the past three decades. Despite the increasing campaign on malaria and the need for protection against infection, malaria prevalence happens to be on the rise in the study area. However, our findings revealed a low prevalence of malaria among the key HIV population in the same study area. This finding serves to highlight the fact that current interventions seem to be yielding little or no results in reducing the incidence of the disease.

4. CONCLUSION

This study further confirmed the presence of HIV/Malaria coinfection in old Cross River State, Nigeria. Our finding has the potential to contribute to knowledge on the epidemiology of HIV/Malaria coinfection. The malaria prevalence reported in this study group differs greatly from what has been reported previously in the general population in the study area. Despite the low coinfection rate obtained in this study, HIV patients seem to be at higher risk of malaria owing to the suppressed immune system and this knowledge calls for more structured approach on the care of those living with HIV. There is still the need for intensified awareness of Malaria and HIV prevention. Our findings also highlight the need for a well-structured approach to the management of HIV/Malaria coinfection in Nigeria.

CONSENT

All authors declare that 'informed consent was obtained from the patient (or other approved

parties) for this study. Consenting HIV-infected individuals accessing HIV clinics at the two tertiary hospitals in old Cross River State, Nigeria who were not on medications for any conditions, were included in the study irrespective of age. Individuals who failed to give their consent were excluded from the study.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed following the ethical standards laid down in the 1964 Declaration of Helsinki." Ethical approval for this study was obtained from the Hospital Research Ethics Committees of Uyo Teaching Hospital (UUTH), Uyo, Nigeria and University of Calabar Teaching Hospital (UCTH), Calabar, Nigeria.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. KM, Williams BG. Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: A mathematical model. *Lancet*. 2009;13:48–57.
2. Granich R, Crowley S, Vitoria M, Lo YR, Souteyrand Y, Dye C, Gilks C, Guerma T, De Cock KM, Williams B. Highly active antiretroviral treatment for the prevention of HIV transmission. *J Int AIDS Soc*. 2010; 13:1.
3. Eyong CA, Kechia FA, Tembe FE, Tatang CA, Wandum GC, Agbor MA, Mbanya DS, Fokunang CN. Evaluation of malaria parasitemia among HIV/AIDS individuals attending the Bamenda regional hospital treatment center. *Central African Journal of Public Health*. 2019;5(2):65-76.
4. World Health Organization. Global malaria control and elimination: report of a technical review. World Health Organization, Geneva. 2008; 47p. Available:<https://apps.who.int/iris/handle/10665/43903> Accessed January 27, 2020.
5. World Health Organization. World malaria report; 2019. Available:<https://www.who.int/malaria/publications/world-malaria-report-2019/en/> Accessed February 7, 2020.

6. Onyenekwe CC, Ukibe N, Meludu SC, Ilika A, Aboh N, Ofiaeli N, Onochie A. Prevalence of malaria as coinfection in HIV-infected individuals in a malaria-endemic area of southeastern Nigeria. *Journal of Vector-Borne Diseases*. 2007; 44(4):250.
7. Audu RA, Onwujekwe DI, Onubogu, CC, Adedoyin JA, Onyejebu N, Mafe AG, Musa AZ. Possible impact of coinfections of tuberculosis and malaria on the CD4+ cell counts of HIV patients in Nigeria. *Annals of African Medicine*. 2005;4(1):10-13.
8. Hochman S, Kim K. The impact of HIV and malaria coinfection: What is known and suggested venues for further study. *Interdisciplinary Perspectives on Infectious Diseases*; 2009.
9. Alemu A, Shiferaw Y, Addis Z, Mathewos B, Birhan W. Effect of malaria on HIV/AIDS transmission and progression. *Parasites and Vectors*. 2013;6(1):18.
10. Kalyesubula I, Musoke-Mudido P, Marum L, et al. Effects of malaria infection in human immunodeficiency virus type-1 infected Ugandan children. *Pediatr Infect Dis. J*. 1997;16:876–881.
11. Kanya MR, Gasasira AF, Yeka A, Bakyaite N, Nsohya SL, Francis D, Havlir D. Effect of HIV-1 infection on antimalarial treatment outcomes in Uganda: A population-based study. *The Journal of Infectious Diseases*. 2006;193(1):9-15.
12. Cohen C, Karstaedt A, Frean J, Thomas J, Govender N, Prentice E, Crewe-Brown H. Increased prevalence of severe malaria in HIV-infected adults in South Africa. *Clinical Infectious Diseases*. 2005;41(11):1631-1637.
13. Saracino A, Nacarapa EA, da Costa Massinga ÉA, Martinelli D, Scacchetti M, de Oliveira C, Macome CA. Prevalence and clinical features of HIV and malaria coinfection in hospitalized adults in Beira, Mozambique. *Malaria Journal*. 2012;11(1): 241.
14. Whitworth J, Morgan D, Quigley M, Smith A, Mayanja B, Eotu H, Omoding N, Okongo M, Malamba S, Ojwiya A. Effect of HIV-1 and increasing immunosuppression on malaria parasitaemia and clinical episodes in adults in rural Uganda: A cohort study. *The Lancet*. 2000;356(9235): 1051-1056.
15. Perrault SD, Hajek J, Zhong K, Owino SO, Sichangi M, Smith, G, et al. Human immunodeficiency virus co-infection increases placental parasite density and transplacental malaria transmission in Western Kenya. *The American Journal of Tropical Medicine and Hygiene*. 2009; 80(1):119-125.
16. Patnaik P, Jere CS, Miller WC, Hoffman IF, Wirima J, Pendame R, Meshnick SR, Taylor TE, Molyneux ME, Kublin JG. Effects of HIV-1 serostatus, HIV-1 RNA concentration and CD4 cell count on the incidence of malaria infection in a cohort of adults in rural Malawi. *J Infect Dis*. 2005; 192:984-991
17. Mouala C, Guiguet M, Houzé S, Damond F, Pialoux G, Viget N, Costagliola D, Le Bras J, Matheron S, FHDH-ANRS CO4 Clinical Epidemiology Group. Impact of HIV infection on severity of imported malaria is restricted to patients with CD4 cell counts < 350 cells/microl. *AIDS*. 2009; 23:1997-2004.
18. Davenport GC, Ouma C, Hittner JB, Were T, Ouma Y, Ong'echa JM, Perkins DJ. Hematological predictors of increased severe anemia in Kenyan children coinfecting with *Plasmodium falciparum* and HIV-1. *American Journal of Hematology*. 2010;85:227-233.
19. Okonko I, Osadebe A, Okoli E, Eke E. Malaria and HIV seropositivity: Study on selected individuals at a tertiary healthcare centre in Port Harcourt, Nigeria. *Microbiologia Medica*. 2019;34(8160):32-36. Available:https://doi.org/10.4081/mm.2019.8160
20. Innocent-Adiele HC. Molecular diversity of HIV-1 subtypes, coinfections, Co-receptor usage and Drug-Resistance Mutations (DRMs) among residents of old Cross River State, Nigeria. A PhD Thesis in the Department of Applied Microbiology, Ebonyi State University, Abakaliki, Nigeria. 2019;253.
21. Akwa Ibom State Government official Website gets a new look – The Premium Herald". *The Premium Herald*. Retrieved 2016-04-07.
22. Nwabueze BO. A constitutional history of Nigeria. C. Hurst and Co LTD, UK. 1982; ISBN 9780905838793.
23. Andem AB, Udofia UU, Okorafor KA, George UU. Bioaccumulation of some heavy metals and Total Hydrocarbon (THC) in the tissues of periwinkle (*Tympanotonus Fuscatus* Var *Radula*) in the Intertidal Regions of Qua Iboe River

- Basin, Ibeno, Akwa Ibom State, Nigeria. *Greener Journal of Biological Sciences*. 2013;3(7):258–264.
24. Reclus É. *Africa*. 3 (Reprint Ed.). Forgotten Books. 1892;341:ISBN 1-4400-9130-7.
 25. Macfarlane SB. Conducting a descriptive survey: 2. Choosing a Sampling Strategy. *Trop. Doctor*. 1997;27(1):14-21.
 26. United Nations AIDS Programme, UNAIDS Global Statistics, 2030 -Ending the AIDS epidemic. The Joint United Nations Programme on HIV/AIDS (UNAIDS) Fact Sheet. 2016;1-16.
Available:https://www.unaids.org/sites/default/files/media_asset/20150901_FactSheet_2015_en.pdf
Accessed August 2019.
 27. Niang L, Winn T, Rusli BN. Practical issues in calculating the sample size for prevalence. *Studies Archives of Orofacial Sciences*. 2006;1:9–14.
 28. Inah SA, Ejemot-Nwadiaro R, Inah JA, Eko JE. Prevalence of malaria among pregnant women and children under five years in Abi Local Government Area, Cross River State, Nigeria. *Asian Journal of Medicine and Health*. 2017;7(1):1-7.
 29. Murray CJ, Ortblad KF, Guinovart C, Lim SS, Wolock TM, Roberts DA, et al. Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990–2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014;384:1005–1070.
 30. Dawaki S, Al-Mekhlafi HM, Ithoi I, Ibrahim J, Atroosh WM, Abdulsalam AM, Sady H, Elyana FN, Adamu AU, Yelwa SI, Ahmed A, Al-Areeqi MA, Subramaniam LR, Nasr NA, Lau Y. Is Nigeria winning the battle against malaria? Prevalence, risk factors and KAP assessment among Hausa communities in Kano State. *Malaria Journal*. 2016;15:351.
 31. Wogu MN, Nduka FO, Wogu MD. Prevalence of malaria parasite infection among pregnant women attending antenatal clinics in Port Harcourt, Rivers State, Nigeria. *International Journal of Tropical Disease and Health*. 2013;3(2): 126-132.
 32. Odikamnor O, Iganga A, Ozowara LN, Okoh N. Prevalence of malaria among pregnant mothers and possible relationship to parity in Abakaliki, Southeast Nigeria. *European Journal of Experimental Biology*. 2014;4(4):15-19.
 33. Raimi OG, Kanu CP. The prevalence of malaria infection in pregnant women living in a suburb of Lagos. Nigeria. *African Journal of Biochemistry Research*. 2010; 4(10):243-245.
 34. Gunn JKL, Ehiri JE, Jacobs ET, Ernest KC, Pettygrove S, Kohler LN, Haenchen SD, Obiefune MC, Ezeanolue CO, Ogidi AG, Ezeanolue EE. Population-based prevalence of malaria among pregnant women in Enugu State, Nigeria: The Healthy Beginning Initiative. *Malaria Journal*. 2015;14(438).
 35. Okoli C, Solomon M. Prevalence of hospital-based malaria among children in Jos, North Central Nigeria *British Journal of Medicine & Medical Research*. 2014; 4(17):3231-3237.
 36. Nwaorgu OC, Orajaka BN. Prevalence of malaria among children 1 – 10 years old in communities in Awka North Local Government Area, Anambra State South East Nigeria. *International Multidisciplinary Journal, Ethiopia*. 2011;5(5):264-281.
 37. Nmadu PM, Peter E, Alexander P, Koggie AZ, Maikenti JI. The prevalence of malaria in children between the ages 2-15 visiting Gwarinpa General hospital life-camp, Abuja, Nigeria. *Journal of Health Science*. 2015; 5(3):47-51.
 38. Olasehinde GI, Ajayi AA, Taiwo SO, Adekeye BT, Adeyeba OA. Prevalence and management of falciparum malaria among infants and children in Ota, Ogun State, Southwestern Nigeria. *African Journal of Clinical and Experimental Microbiology*. 2010;11(3):159-163.
 39. Otu AA, Udoh UA, Ita OI, Hicks JP, Ukpeh I, Walley J. Prevalence of Zika and malaria in patients with fever in secondary healthcare facilities in South-Eastern Nigeria. *Trop Doct*. 2020;50(1):22-30.
 40. Orok DA, Ibor UA, Oyama IO, Duke Edisua E, Efeffiom EE, Ekup EU, Unimke A, Akung UJ. Prevalence of malaria and typhoid fever co-infection among febrile patients attending College of Health Technology Medical Centre in Calabar, Cross River State, Nigeria. *International Journal of Current Microbiology and Applied Sciences*. 2016;5:825-835.
 41. Ibor UW, Okoronkwo EM, Rotimi EM. Temporal analysis of malaria prevalence in Cross River State, Nigeria. *E3 Journal of Medical Research*. 2016;5(1):001-007.
 42. Ibor UW, Aigbe GO, Iwara AI, Okongor OE, Okino II. Ownership and utilization of

- insecticide-treated nets in Cross River State, Nigeria. *Journal of Medical Sciences*. 2012;12:198-206.
43. Dada EO, Okebugwu QC, Ibukunoluwa MR. Coinfection of Human Immunodeficiency Virus (HIV) with malaria in Gbalegi, Idanre and State Hospital, Akure, Ondo State, Nigeria. *HIV Current Research*. 2016;1(111):2.
44. Olusi TA, Abe AF. Coinfection of HIV and malaria parasites in pregnant women attending major ante-natal health facilities in Akure, Ondo State, Nigeria. *Journal of Parasitology and Vector Biology*. 2014; 6(9):124-130.
45. Njunda LA, Kamga HLF, Nsagha DS, Assob JCN, Kwenti TE. Low malaria prevalence in HIV-positive patients in Bamenda, Cameroon. *Journal of Microbiology Research*. 2012;2(3):56-59.
46. Jenkins R, Omollo R, Ongecha M, Sifuna P, Othieno C, Onger L, Kingora J, Ogutu B. Prevalence of malaria parasites in adults and its determinants in malaria-endemic area of Kisumu County, Kenya. *Malaria Journal*. 2015;14: 263.
47. Bhattacharya MK, Naik TN, Ghosh M, Jana S, Dutta P. Pulmonary tuberculosis among HIV seropositives attending a counselling center in Kolkata. *Indian Journal of Public Health*. 2011;55(4):329.
48. United States Embassy in Nigeria. Nigeria Malaria Fact Sheet. Economic Section, United States Embassy in Nigeria Plot 1075, Diplomatic Drive Central Area Abuja, FCT, Nigeria. Pp.2. Available: <https://photos.state.gov/libraries/nigeria/231771/Public/December-MalariaFactSheet2.pdf> Accessed July 14, 2020
49. World Health Organization. World Malaria Report, 2013. World Health Organization, Geneva. 2013;284. Available: https://www.who.int/malaria/publications/world_malaria_report_2013/en/ Accessed January 27, 2020.
50. Kublin JG, Patnaik P, Jere CS, Miller WC, Hoffman IF, Chimbiya N, Pendame R, Taylor TE, Molyneux ME. Effect of *Plasmodium falciparum* malaria on concentration of HIV-1-RNA in the blood of adults in rural Malawi: A prospective cohort study. *Lancet*. 2005;365: 33-240.

© 2020 Ejike et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/57414>