



## Effect of Brassinolide on the Growth of Mustard Crops Grown in Semi-arid Tropics of Nizamabad

P. Latha<sup>1</sup> and B. Vidya Vardhini<sup>2\*</sup>

<sup>1</sup>Department of Botany, Govt. Degree College, Armoor, Nizamabad, India.

<sup>2</sup>Department of Botany, Telangana University, Dichpally, Nizamabad -503322, India.

### Authors' contributions

This work was carried out in collaboration between both authors. Author BVV is the research supervisor and author PL is pursuing her Ph. D under author BVV. Author BVV designed the study, wrote the protocol and wrote the first draft of the manuscript. Author PL managed the literature searches; analyses of the study performed the spectroscopy analysis and managed the experimental process. Both the authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/IJPSS/2016/21617

#### Editor(s):

(1) A. Mujib, Department of Botany, Hamdard University, India.

#### Reviewers:

(1) Geetika Sirhindi, Punjabi University, India.

(2) Keimei Oh, Akita Prefectural University, Japan.

(3) Mayank Anand Guruani, Yeungnam University, South Korea.

(4) Mohammed Suleiman, Umaru Musa Yar'adua University, Nigeria.

(5) Ndamuleleni Romeo Murovhi, Institute for Tropical and Sub-tropical Crops, South Africa.

Complete Peer review History: <http://sciencedomain.org/review-history/11926>

Original Research Article

Received 26<sup>th</sup> August 2015

Accepted 1<sup>st</sup> October 2015

Published 20<sup>th</sup> October 2015

### ABSTRACT

The effect of Brassinolide (BL) sprayed in three concentrations viz., 0.5  $\mu$ M, 1.0  $\mu$ M and 2.0  $\mu$ M on the shoot growth, root growth and foliar growth (plant fresh weight, leaves per plant and leaf area) of mustard plants grown in the semi-arid tropics of Nizamabad was studied. The soil in Nizamabad district is saline land black soil wherein the plants usually experience drought and saline stresses. All the three concentrations of BL stimulated the shoot growth, root growth as well as the foliar growth of mustard plants. BL at 0.5 $\mu$ M was found most effective in increasing the shoot and root growth. The promotion of shoot, root and foliar growth is an indicator that BL mitigated the negative effect of the semi-arid conditions of the soil.

**Keywords:** Brassinolide; foliar growth; mustard; root growth; shoot growth.

\*Corresponding author: E-mail: [dvidyavardhini@rediffmail.com](mailto:dvidyavardhini@rediffmail.com);

## ABBREVIATIONS

*Brassinolide=BL; Brassinosteroids =BRs; 24-Epibrassinolide = 24-EpiBL; 28-Homobrassinolide =28-Homo BL.*

## 1. INTRODUCTION

Brassinosteroids (BRs) are a new type of polyhydroxy steroidal phytohormones with significant growth-promoting influence [1]. Mitchell et al. [2] discovered BRs which were later extracted from the pollen of *Brassica napus* L. by Grove et al. [3]. BRs can be classified as C<sub>27</sub>, C<sub>28</sub> or C<sub>29</sub> BRs according to the number of carbons in their structure [4]. However, Vardhini et al. [5] reported that brassinolide (BL), 28-homobrassinolide (28-HomoBL) and 24-epibrassinolide (24-EpiBL) are the three bioactive BRs being widely used in most physiological and experimental studies. The work with BR biosynthetic mutants in *Arabidopsis thaliana* [6] and *Pisum sativum* [7] have provided strong evidences that BRs are essential for plant growth and development and BR- signaling plays a positive in plant growth and development viz., spatiotemporal control of BR pathways in plant development employing microscope lens turret to study the pleiotropic phenotypes of the BR mutants at a higher magnification [8].

Rao et al. [9] stated that BRs are a new group of plant growth hormones that perform a variety of physiological roles like growth, seed germination, rhizogenesis, senescence, etc. and also confer resistance to plants against various abiotic stresses. Though, BRs were initially identified based on their growth promoting activities, subsequent physiological and genetic studies revealed additional functions of BRs in regulating a wide range of processes, including source/sink relationships, seed germination, photosynthesis, senescence, photomorphogenesis, flowering and responses to different abiotic and biotic stresses [4].

Nizamabad district experiences a tropical dry and wet season with most of the rainfall in June to October. It usually experiences erratic rain fall. The soil is saline land black soil which is deep loamy to clay loam, moderately drained, neutral to alkaline in nature. The reduction of growth of many plants by salinity and drought usually effects on dry matter production, ionic relations, metabolic variations, physiological processes, water contents. The semi-arid condition directly poses a threat to the overall yield of the plants as

they usually experience drought and saline stresses.

Mustard (*Brassica juncea*) is an oil yielding plant that belongs to the family Brassicaceae which is grown throughout the world. The plant has tiny seeds which are usually used as a condiment. Mustard oil is extracted from the seed which are used for cooking, massaging etc. Some kinds of mustard plants have edible leaves that are known as *mustard greens*. It is a well-established fact from time immemorial that plants are the vital components of dietary food chains in which they provide almost all the essential mineral and organic nutrients to humans [10]. The present study is undertaken to understand the effect of application of BL on the growth of mustard plants in terms of shoot, root and foliage grown in the semi - arid soils of Nizamabad.

## 2. MATERIALS AND METHODS

### 2.1 Chemicals and Plant Material

Brassinolide (BL: double) is a commercially available brassinosteroid was procured from Bahar Agrochem & Feeds Pvt. Ltd, Ratnagiri, Maharashtra, India, Ltd. It is marketed by Godrej Agrovet Ltd., Hyderabad, Andhra Pradesh, India. Brassinolide (Double) consists of 0.1% of brassinolide, 2.0% of emulsifier and 97.9% of solvent IPA.

Mustard (*Brassica juncea* var tulasi) was purchased from National Seeds Corporation, Hyderabad, Andhra Pradesh, India.

### 2.2 Shoot and Root Growth

Mustard seeds were sown in clay pots containing fresh sieved black soil mixed with farmyard manure. Plants were grown under natural day length. BL was supplied to the plants as foliar spray at three different concentration levels viz., 0.5  $\mu$ M, 1.0  $\mu$ M and 2.0  $\mu$ M on 35<sup>th</sup>, 40<sup>th</sup>, 45<sup>th</sup> and 50<sup>th</sup> day (from the day of sowing). Growth parameters in terms of shoot weight diameter and length were recorded on the 55<sup>th</sup> day. Growth parameters in terms of root weight, diameter and length were recorded on the 55<sup>th</sup> day.

### 2.3 Foliar Growth

Foliar growth was recorded in terms of plant fresh weight, leaves per plant and leaf area per plant were recorded on 55<sup>th</sup> day. Average leaf area per plant was determined with the help of the equation developed by Kemp [11].

$$A = (L \times B \times 0.9) \times n,$$

Where A= leaf area, L= leaf length, B=leaf breadth, 0.9= constant factor and n= number of leaves.

The data is represented in terms of Mean±S.E (n=9).

### 3. RESULTS

Exogenous application of BL resulted in substantial increase in shoot growth of mustard plants as reflected in increases in length, fresh weight and diameter of the shoots (Table 1.). All the three concentrations applied viz., 0.5µM, 1 µM and 2 µM increased the shoot growth of mustard plants grown in semi-arid soils of Nizamabad over control plants. BL at 0.5µM was found most effective in substantial increase in

shoot growth compared to the other two concentrations as well as control plants.

The mustard plants grown in semi-arid soils of Nizamabad and treated with foliar application of BL showed increased root growth in terms of length, fresh weight and diameter (Table 2.). All the three concentrations of BL applied viz., 0.5 µM, 1 µM and 2 µM increased the shoot growth of mustard plants grown in semi-arid soils of Nizamabad over control plants. 0.5 µM Conc. of BL exhibited maximum enhanced root growth compared to the other treatments as well as untreated controls.

Foliar application of BL exhibited improved foliage in terms of plant fresh weight, leaves/plant and leaf area of mustard plants grown in semi-arid tropics of Nizamabad (Table 3.). All the three concentrations applied viz., 0.5 µM, 1 µM and 2 µM increased the shoot growth of mustard plants grown in semi-arid soils of Nizamabad over control plants. Application of BL at 0.5 µM conc. was more pronounced in increasing the foliar growth of mustard plants compared to the other two treatments as well as control plants.

**Table 1. Effect of brassinolide on the shoot growth in *Brassica juncea***

Treatment (µM)	Shoot weight (gm/fr. wt.)*	Shoot diameter (cm)*	Shoot length (cm)*
0.5	8.093±1.925	6.400±0.213	61.66±3.179
1.0	6.610±0.100	6.427±0.019	57.10±1.000
2.0	4.510±0.238	5.098±0.036	47.66±7.881
Control	2.001±1.041	4.330±0.169	32.00±4.582

\*Values are Mean±S.E. (N=9), (gm/fr. wt.) = gram/fresh weight; (cm) = centimeters

**Table 2. Effect of brassinolide on the root growth in *Brassica juncea***

Treatment (µM)	Root weight (gm/fr. wt.)*	Root diameter (cm)*	Root length (cm)*
0.5	1.130±0.430	4.850±0.087	11.02±0.288
1.0	0.830±0.100	4.370±0.134	10.50±0.288
2.0	0.710±0.058	4.435±0.131	10.33±0.033
Control	0.440±0.145	3.320±0.150	8.83±4.582

\*Values are Mean±S.E. (N=9), (gm/fr. wt.) = gram/fresh weight; (cm) = centimeters

**Table 3. Effect of brassinolide on the foliar growth of *Brassica juncea***

Treatment (µM)	Plant fresh weight (gm/fr. wt.)*	Number of leaves/plant	Leaf area (sq. cm)*
0.5	9.550±2.224	12.33±1.850	483.1±11.26
1.0	8.403±1.315	11.26±1.650	411.3±9.90
2.0	5.510±1.043	10.66±2.060	398.2±10.10
Control	2.991±1.041	7.33±1.520	133.1±9.91

\*Values are Mean±S.E. (N=9), (gm/fr. wt.) = gram/fresh weight; (sq cm) = square centimeters

#### 4. DISCUSSION

It is a well-established fact that BRs are a new group of plant growth regulators which play a positive role in the growth and development of plants. BRs increased the height of two field-grown inbred lines of maize (*Zea mays* L.) during the vegetative and early reproductive phases of plant development during the early weeks after their application [12]. Fariduddin et al. [13] reported that cucumber seedlings sprayed with  $10^{-8}$ , or  $10^{-6}$  M of 28-homoBL exhibited improved growth at the 30-day stage after treatment plants were exposed for 18 h to chilling temperature (10/8°C, 5/3°C). BL applied to the sunflower seedlings significantly increased the hypocotyl growth under different light qualities [14]. Even in the present study, application of BL significantly increased the shoot growth in mustard plants grown in the semi – arid soils of Nizamabad.

Vardhini et al. [15] reported that application of BRs increased the qualitative growth of radish (*Raphanus sativus* L. var Pusa chetki long) in terms of root fresh weight, root dry weight, oxalic acid, ascorbic acid, folic acid and niacin. Hayat et al. [16] reported that 28-homoBL and 24-epiBL increased the growth of *Lycopersicon esculentum*. Imbibition of 24-epiBL to pea (*P. sativum* L.) cv. Climax seeds subjected to sodium chloride stress exhibited increased germination, embryo axis length and most of the aspects of shoot and root growth at seedling stage [17]. Earlier studies clearly emphasized that external supplementation/application of BRs improved plant growth in the case of wheat [18] and groundnut [19] which is in tune with the present study where foliar application of BL substantially increased the root growth of mustard plants.

The enhanced shoot and root growth of BL treated mustard plants was associated with increased foliar growth. Arteca and Arteca [20] also reported that BRs not only induce exaggerated growth in hydroponically grown *A. thaliana*, but also control the proliferation of its leaf cells [21]. Kudryashova et al. [22] reported that supplementation of 24-epiBL played a pivotal role in *in vitro* regeneration of highbush blueberry (*Vaccinium corymbosum* L., cv. Brigitta blue).

#### 5. CONCLUSION

The present study reveals that application of BL on mustard plants as foliar spray promoted the

shoot, root and foliar growth of mustard plants grown in semi-arid soils of Nizamabad. The soils of Nizamabad are saline and dry in nature inhibiting the growth of plants. BRs have the ability to promote growth of plant under stressful conditions. The present study reveals a new insight that application of BL overcame the negative effect of the semi-arid conditions of the soil (reflected in the control plants) and promoted the shoot, root and foliar growth (reflected in the BL-treated plants) of mustard.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Vardhini BV, Anjum NA. Brassinosteroids make plant life easier under abiotic stresses mainly by modulating major components of antioxidant defense system. *Frontiers in Environmental Sciences*. 2015;2:67. DOI: 10.3389/fenvs.2014.0006.
2. Mitchell JW, Mandava NB, Worley JE, Plimmer JR, Smith MV. Brassins: A family of plant hormones from rape pollen. *Nature*. 1970;225:1065-1066.
3. Grove MD, Spencer GF, Rohwedder WK, Mandava NB, Worlet JF, Warthen Jr JC, Steffens GL, Flippen-Andersen JL, Cook Jr JC. Brassinolide, a plant promoting steroid isolated from *Brassica napus* pollen. *Nature*. 1979;281:121-124.
4. Vardhini BV. Brassinosteroids, Role for amino acids, peptides and amines modulation in stressed plants - A review. In: Anjum NA, Gill SS, Gill R, editors. *Plant Adaptation to Environmental Change: Significance of Amino Acids and their Derivatives*, United Kingdom: CAB International of Nosworthy Way, Wallingford OX10 8DE. 300-316; 2013.
5. Vardhini BV, Anuradha S, Rao SSR. Brassinosteroids - New class of plant hormone with potential to improve crop productivity. *Indian J. Plant Physiol*. 2006;11:1-12.
6. Tao YZ, Zheng J, Xu ZM, Zhang XH, Zhang K, Wang GY. Functional analysis of ZmDWF1, a maize homolog of *Arabidopsis* brassinosteroids biosynthetic DWF1/DIM gene. *Plant Sci*. 2004;167:743-751.
7. Nomura T, Nakayama N, Reid JB, Takeuchi Y, Yokota T. Blockage of

- brassinosteroid biosynthesis and sensitivity cause dwarfism in *Pisum sativum*. Plant Physiol. 1997;113:31-37.
8. Fábregas N, Caño-Delgado AI. Turning on the microscope turret: A new view for the study of brassinosteroid signaling in plant development. Physiol Plant. 2014;151:172-183.
  9. Rao SSR, Vardhini BV, Sujatha E, Anuradha S. Brassinosteroids – new class of phytohormones. Curr Sci. 2002;82: 1239-1245.
  10. Grusak MA, Dellapenna D. Improving the nutrient composition of plants to enhance human nutrition and health. Annu Rev Plant Physiol Plant Mol Biol. 1999;50:133-161.
  11. Kemp CD. Methods of estimating leaf area of grasses from linear measurement. Ann Bot. 1960;24:491-512.
  12. Hola D, Rothova O, Kocova M, Kohout L, Kvasnica M. The effect of brassinosteroids on the morphology, development and yield of field-grown maize. Plant Growth Regul. 2010;61:29-43.
  13. Fariduddin Q, Yusuf M, Chalkoo S, Hayat S, Ahmad A. 28- Homobrassinolide improves growth and photosynthesis in *Cucumis sativus* L. through an enhanced antioxidant system in the presence of chilling stress. Photosynthetica. 2011;49:55-64.
  14. Kurepin LV, Joo SH, Kim SK, Pharis RP, Back TG. Interaction of brassinosteroids with light quality and plant hormones in regulating shoot growth of young sunflower and *Arabidopsis* seedlings. J Plant Growth Regul. 2012;31:156-164.
  15. Vardhini BV, Sujatha E, Rao SSR. Studies on the effect of brassinosteroids on the qualitative changes in the storage roots of radish. Bulg J Agric Sci. 2012;18:63-69.
  16. Hayat S, Yadav S, Wani AS, Irfan M, Ahmad A. Comparative effect of 28-homobrassinolide and 24-epibrassinolide on the growth, carbonic anhydrase activity and photosynthetic efficiency of *Lycopersicon esculentum*. Photosynthetica. 2011;49:397-404.
  17. Shahid MA, Pervez MA, Balal RM, Mattson NS, Rashid A, Ahmad R, Ayyub CM, Abbas T. Brassinosteroid (24-epibrassinolide) enhances growth and alleviates the deleterious effects induced by salt stress in pea (*Pisum sativum* L.). Aust J Crop Sci. 2011;5:500-510.
  18. Sairam RK. Effect of homobrassinolide application on plant metabolism and grain yield under irrigated and moisture – stress conditions of two wheat varieties. Plant Growth Regul. 1994;14:173-181.
  19. Vardhini BV, Rao SSR. Effect of brassinosteroids on growth, metabolite content and yield of *Arachis hypogaea*. Phytochemistry. 1998;48:927-930.
  20. Arteca JM, Arteca RN. Brassinosteroid-induced exaggerated growth in hydroponically grown *Arabidopsis* plants. Physiol Plant. 2001;112:104-112.
  21. Nakaya M, Tsukaya H, Murakami N, Kato M. Brassinosteroids control the proliferation of leaf cells of *Arabidopsis thaliana*. Plant Cell Physiol. 2002;43:239-244.
  22. Kudryashova OA, Volotovich AA, Vasilevskaya TI, Varavina NP, Rupasova ZA, Khripach VA. Effects of 24-epibrassinolide on in vitro micropropagation of highbush blueberry. Russ J Plant Physiol. 2012;59:586-593.

© 2016 Latha and Vardhini; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://sciencedomain.org/review-history/11926>